

5

10


15

Found : C, 78.40; H, 7.09; N, 6.09.

Melting point: 179 - 181°C (crystallization solvent: ethyl acetate)

20

20



25

¹H NMR (CDCl₃) δ: 1.25-1.71 (8H, m), 1.95-2.00 (2H, m), 2.25-2.45 (6H, m), 2.83-2.93 (3H, m), 7.09 (1H, d, J = 8.3 Hz), 7.30-7.32 (1H, m), 7.43- 7.45 (3H, m), 7.55 (2H, d, J = 8.1 Hz), 7.65 (2H, d, J = 8.4 Hz), 7.77 (1H, s), 7.93 (2H, d, J = 8.1 Hz).

WO 01/21577


PCT/JP00/06375

201

Melting point: 202 - 203°C (crystallization solvent:
tetrahydrofuran - n-hexane)

Example 56

5 5-Oxo-1-phenyl-N-[6-(1-piperidinylmethyl)-5,6,7,8-
tetrahydro-2-naphthalenyl]-3-pyrrolidinecarboxamide



The titled compound was obtained by carrying out the
same operation as in Example 1, using 6-(1-
10 piperidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine
obtained in Reference Example 53.

^1H NMR (CDCl_3) δ : 1.03-3.33(22H, m), 3.97 (1H, t, J = 8.4
Hz), 4.21 (1H, dd, J = 6.8, 7.1 Hz), 6.91-7.63 (9H, m).

Elemental analysis for $\text{C}_{27}\text{H}_{33}\text{N}_3\text{O}_2$


15 Calcd.: C, 75.14; H, 7.71; N, 9.74.

Found: C, 75.01; H, 7.33; N, 9.43.

Melting point: 162 - 164°C (crystallization solvent: ethyl
acetate)

20 Example 57

6-(4-Chlorophenyl)-N-[6-(1-piperidinylmethyl)-5,6,7,8-
tetrahydro-2-naphthalenyl]nicotinamide



The titled compound was obtained by carrying out the
same operation as in Example 1, using 6-(1-
25 piperidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine
obtained in Reference Example 53.

^1H NMR (CDCl_3) δ : 1.30-2.40 (16H, m), 2.82-2.92 (3H, m),
7.09 (1H, d, J = 8.1 Hz), 7.26-7.48 (4H, m), 7.80 (2H, d,

WO 01/21577

PCT/JP00/06375

202


$J = 8.7 \text{ Hz}$), 7.99 (2H, d, $J = 8.7 \text{ Hz}$), 8.23 (d, 1H, $J = 6.3 \text{ Hz}$), 9.11 (1H, s).

Melting point: 193 - 195°C (crystallization solvent: ethyl acetate)

5

Example 58

5-(4-Chlorophenyl)-N-[6-(1-piperidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenyl]-2-furamide




10 The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-piperidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine obtained in Reference Example 53.

$^1\text{H NMR}$ (CDCl_3) δ : 1.23-1.61 (7H, m), 1.96-2.00 (2H, m),
15 2.24-2.43 (7H, m), 2.80-2.92 (3H, m), 6.75 (1H, d, $J = 3.6 \text{ Hz}$), 7.07 (1H, d, $J = 8.4 \text{ Hz}$), 7.27 (1H, d, $J = 3.6 \text{ Hz}$), 7.32-7.42 (4H, m), 7.66 (2H, d, $J = 8.4 \text{ Hz}$), 8.32 (1H, s).

Example 59

20 N-[6-(1-Piperidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenyl]-3-(2,4,5-triethoxyphenyl)-5-isoxazolecarboxamide



25 The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-piperidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine obtained in Reference Example 53.

$^1\text{H NMR}$ (CDCl_3) δ : 1.42-1.60 (18H, m), 1.97-2.36 (7H, m),
2.80-2.95 (3H, m), 4.06-4.18 (6H, m), 6.58 (1H, s), 7.09
30 (1H, d, $J = 8.4 \text{ Hz}$), 7.35 (1H, d, $J = 8.1 \text{ Hz}$), 7.44 (1H, s), 7.50 (1H, s), 7.55 (1H, s), 8.16 (1H, s).


WO 01/21577

PCT/JP00/06375

203

Example 60

4-(4-Chlorophenyl)-2-phenyl-N-[6-(1-piperidinylmethyl)-
5,6,7,8-tetrahydro-2-naphthalenyl]-1,3-oxazole-5-
5 carboxamide




The titled compound was obtained by carrying out the
same operation as in Example 1, using 6-(1-
piperidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine
10 obtained in Reference Example 53.

¹H NMR (CDCl₃) δ: 1.26-1.58 (7H, m), 1.90-2.00 (2H, m),
2.22-2.35 (7H, m), 2.70-2.95 (3H, m), 7.06 (1H, d, J = 8.1
Hz), 7.25-7.51 (7H, m), 8.04-8.32 (5H, m).

15 Example 61

4'-Chloro-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-
naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the
20 same operation as in Example 51, using 4'-chloro-N-[6-
(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-
biphenyl]-4-carboxamide obtained in Reference Example 56.
Melting point: 185 - 187°C (crystallization solvent:
tetrahydrofuran - n-hexane)

25 ¹H NMR (CDCl₃) δ: 1.83 (4H, s), 2.35 (2H, t, J = 8.1 Hz),
2.52 (4H, s), 2.84 (2H, t, J = 8.1 Hz), 3.18 (2H, s), 6.36
(1H, s), 7.02 (1H, d, J = 8.4 Hz), 7.39-7.56 (6H, m), 7.66

WO 01/21577


PCT/JP00/06375

204

(2H, d, J = 7.5 Hz), 7.82 (1H, s), 7.93 (2H, d, J = 7.5 Hz).

Example 62


5 5-(4-Chlorophenyl)-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]-2-pyridinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 51, using 4'-chloro-N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 56.
¹H NMR (CDCl₃) δ: 1.80 (6H, s), 2.37 (2H, t, J = 8.1 Hz), 2.52 (4H, s), 2.87 (2H, t, J = 8.1 Hz), 3.18 (2H, s), 6.37 (1H, s), 7.03 (1H, d, J = 7.8 Hz), 7.48-7.61 (6H, m), 8.04 (1H, dd, J = 8.1, 2.1 Hz), 8.35 (1H, d, J = 8.1 Hz), 8.78 (1H, s), 9.95 (1H, s).

Example 63

4-(4-Pyridinyl)-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]benzamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.
¹H NMR (CDCl₃) δ: 1.79-1.83 (6H, m), 2.35 (2H, t, J = 8.1 Hz), 2.53 (4H, s), 2.73 (2H, t, J = 8.1 Hz), 3.18 (2H, s), 6.36 (1H, s), 7.02 (1H, d, J = 7.8 Hz), 7.38 (1H, d, J = 8.1 Hz), 7.48 (1H, s), 7.71-7.78 (4H, m), 7.89 (1H, s), 7.99 (1H, d, J = 8.4 Hz), 8.32 (2H, d, J = 8.4 Hz).


WO 01/21577

PCT/JP00/06375

205

Example 64

4'-Chloro-N-[6-[(4-phenyl-1-piperidiny)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 51, using 4'-chloro-N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 56.


10 ¹H NMR (CDCl₃) δ: 1.83-2.10 (6H, m), 2.37 (2H, t, J = 8.1 Hz), 2.47-2.54 (1H, m), 2.86 (2H, t, J = 8.1 Hz), 3.03-3.10 (2H, m), 3.10 (2H, s), 6.37 (1H, s), 7.03 (1H, d, J = 8.4 Hz), 7.19-7.57 (11H, m), 7.66 (2H, d, J = 8.4 Hz), 7.81 (1H, s), 7.94 (2H, d, J = 8.4 Hz).

15 Melting point: 228 - 230°C (crystallization solvent: tetrahydrofuran - n-hexane)

Example 65

4'-Chloro-N-[6-(4-morpholinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

20



The titled compound was obtained by carrying out the same operation as in Example 51, using 4'-chloro-N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 56.

25 ¹H NMR (CDCl₃) δ: 2.34 (2H, t, J = 7.8 Hz), 2.45 (4H, s), 2.84 (2H, t, J = 7.8 Hz), 3.06 (2H, s), 3.73 (4H, s), 6.36 (1H, s), 7.02 (1H, d, J = 8.1 Hz), 7.36-7.57 (6H, m), 7.67 (2H, d, J = 8.4 Hz), 7.80 (1H, s), 7.94 (2H, d, J = 8.4 Hz).

WO 01/21577


PCT/JP00/06375

206

Melting point: 194 - 195°C (crystallization solvent:
tetrahydrofuran - n-hexane)

Example 66

5 4'-Chloro-N-(6-[[methyl(2-phenylethyl)amino]methyl]-
7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide




The titled compound was obtained by carrying out the
same operation as in Example 51, using 4'-chloro-N-[6-
10 (chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-
biphenyl]-4-carboxamide obtained in Reference Example 56.
¹H NMR (CDCl₃) δ: 2.25-2.32 (2H, m), 2.32 (3H, s), 2.60-2.66
(2H, m), 2.77-2.83 (4H, m), 3.10 (2H, s), 6.32 (1H, s),
6.93-7.95 (16H, m).

15 Melting point: 173 - 175°C (crystallization solvent:
tetrahydrofuran - n-hexane)

Example 67

20 4'-Chloro-N-[6-[methylanilino)methyl]-7,8-dihydro-2-
naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the
same operation as in Example 51, using 4'-chloro-N-[6-
(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-
25 biphenyl]-4-carboxamide obtained in Reference Example 56.
¹H NMR (CDCl₃) δ: 2.20-2.30 (2H, m), 2.25 (3H, s), 2.85-2.90

WO 01/21577


PCT/JP00/06375

207

(2H, m), 3.00 (2H, s), 6.30 (1H, s), 6.74-7.95 (146H, m).
Melting point: 177 - 179°C (crystallization solvent:
tetrahydrofuran - n-hexane)

5 Example 68


4'-Chloro-N-[6-[(4-phenyl-1-piperadiny)methyl]-7,8-
dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



10 The titled compound was obtained by carrying out the
same operation as in Example 51, using 4'-chloro-N-[6-
(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-
biphenyl]-4-carboxamide obtained in Reference Example 56.
¹H NMR (CDCl₃) δ: 2.37 (2H, t, J = 8.1 Hz), 2.62 (4h, S),
2.86 (2H, t, J = 8.4 Hz), 3.13 (2H, s), 3.22 (4H, s), 6.39
15 (1H, s), 6.85-7.95 (16H, m).
Melting point: 228 - 230°C (crystallization solvent:
tetrahydrofuran - n-hexane)

Example 69

20 4'-Chloro-N-[6-[[[2-
(dimethylamino)ethyl](methyl)amino)methyl]-7,8-dihydro-
2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



25 The titled compound was obtained by carrying out the
same operation as in Example 51, using 4'-chloro-N-[6-
(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-
biphenyl]-4-carboxamide obtained in Reference Example 56.
¹H NMR (CDCl₃) δ: 2.25 (6H, s), 2.26 (3H, s), 2.33 (2H, t,

WO 01/21577

PCT/JP00/06375


208

$J = 8.1$ Hz), 2.44-2.50 (4H, m), 2.84 (2H, t, $J = 8.1$ Hz), 3.07 (2H, s), 6.35 (1H, s), 7.02 (1H, d, $J = 8.4$ Hz), 7.37-7.57 (6H, m), 7.67 (2H, d, $J = 8.1$ Hz), 7.80 (1H, s), 7.94 (2H, d, $J = 8.4$ Hz).

5 Melting point: 156 - 158°C (crystallization solvent: tetrahydrofuran - n-hexane)

Example 70

4'-Fluoro-N-[6-(4-morpholinylmethyl)-5,6,7,8-
10 tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide




The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(4-morpholinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine
15 obtained in Reference Example 57.

^1H NMR (CDCl_3) δ : 1.40-1.50 (1H, m), 1.90-2.10 (2H, m), 2.29-2.45 (7H, m), 2.80-2.92 (3H, m), 3.72-3.75 (4H, m), 7.07-7.33 (4H, m), 7.46 (1H, s), 7.56-7.66 (4H, m), 7.78 (1H, s), 7.92 (2H, d, $J = 8.1$ Hz).

20 Melting point: 188 - 190°C (crystallization solvent: ethyl acetate)

Example 71

4'-Chloro-N-[6-(4-morpholinylmethyl)-5,6,7,8-
25 tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(4-morpholinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine

WO 01/21577

PCT/JP00/06375

209


obtained in Reference Example 57.

¹H NMR (CDCl₃) δ: 1.40-1.50 (1H, m), 1.90-2.10 (2H, m),
2.32-2.45 (7H, m), 2.80-2.90 (3H, m), 3.70-3.80 (4H, m),
7.10-7.92 (12H, m).

5 Melting point: 216 - 218° C (crystallization solvent: ethyl
acetate)

Example 72

10 4-Chloro-N-[6-(4-morpholinylmethyl)-5,6,7,8-tetrahydro-
2-naphthalenyl]-2-phenyl-5-pyrimidinecarboxamide




The titled compound was obtained by carrying out the
same operation as in Example 1, using 6-(4-
morpholinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine
15 obtained in Reference Example 57.

¹H NMR (CDCl₃) δ: 1.40-1.50 (1H, m), 1.95-2.05 (2H, m),
2.29-2.45 (7H, m), 2.80-2.95 (3H, m), 3.73 (4H, t, J = 4.5
Hz), 7.10 (1H, d, J = 8.1 Hz), 7.32 (1H, d, J = 8.1 Hz),
7.42 (1H, s), 7.49-7.56 (3H, m), 8.25 (1H, s), 8.48 (2H,
20 d, J = 6.6 Hz), 9.20 (1H, s)

Example 73

N-[6-(4-Morpholinylmethyl)-5,6,7,8-tetrahydro-2-
naphthalenyl]-2-phenyl-5-pyrimidinecarboxamide



25

The titled compound was obtained by carrying out the
same operation as in Reference Example 48, using 4-
chloro-N-[6-(4-morpholinylmethyl)-5,6,7,8-tetrahydro-2-
naphthalenyl]-2-phenyl-5-pyrimidinecarboxamide obtained in

WO 01/21577

PCT/JP00/06375


210

Example 72.

¹H NMR (CDCl₃) δ: 1.21-1.30 (1H, m), 1.93-2.03 (2H, m), 2.28-2.44 (7H, m), 2.80-2.90 (3H, m), 3.73 (4H, t, J = 4.8 Hz), 7.07 (1H, d, J = 8.1 Hz), 7.26-7.30 (1H, m), 7.39 (1H, s), 7.51-7.53 (3H, m), 8.00 (1H, s), 8.50 (2H, dd, J = 8.1, 2.4 Hz), 9.21 (2H, s)

Example 74


N-[6-[(Diethylamino)methyl]-7,8-dihydro-2-naphthalenyl]
[1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 51, using N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 58.
¹H NMR (CDCl₃) δ: 1.24 (6H, t, J = 7.2 Hz), 2.33 (2H, t, J = 5.1 Hz), 2.53 (4H, q, J = 7.2 Hz), 2.84 (2H, t, J = 5.1 Hz), 3.11 (2H, s), 6.36 (1H, s), 7.02 (1H, d, J = 8.1 Hz), 7.37-7.50 (5H, m), 7.63 (2H, d, J = 8.7 Hz), 7.71 (2H, d, J = 8.4 Hz), 7.79 (1H, s), 7.93 (2H, d, J = 8.4 Hz).
Melting point: 153 - 155°C (crystallization solvent: tetrahydrofuran - n-hexane)

Example 75

4-(2-Benzo[b]furanyl)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]benzamide



The titled compound was obtained by carrying out the

WO 01/21577

PCT/JP00/06375

211


same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

Melting point: 192 - 194°C (crystallization solvent: tetrahydrofuran-isopropyl ether)

5

Example 76

4-(3-Methoxybenzyloxy)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]benzamide



10


The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

Melting point: 102 - 104°C (crystallization solvent: isopropyl ether)

15

Example 77

4-(4-Fluorobenzyloxy)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]benzamide



20

The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

Melting point: 165 - 167°C (crystallization solvent: tetrahydrofuran-hexane)

25


Example 78

4-[4-(Methylsulfonyl)benzyloxy]-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]benzamide

WO 01/21577

PCT/JP00/06375

212




The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

5 Melting point: 162 - 163°C (crystallization solvent: tetrahydrofuran-hexane)

Example 79

10 4-(4-Ethylbenzyloxy)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]benzamide




The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

15 Melting point: 120 - 122°C (crystallization solvent: tetrahydrofuran-isopropyl ether)

Example 80

20 (4'-Methylbiphenyl-4-yl)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]carboxamide



The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-

WO 01/21577

PCT/JP00/06375


213

dimethylamino)methyl]tetralin hydrochloride.

Melting point: 181 - 182°C (crystallization solvent: ethyl acetate-hexane)

5 Example 81

(2',4'-Dichlorobiphenyl-4-yl)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]carboxamide




10 The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

Melting point: 188 - 189°C (crystallization solvent: tetrahydrofuran-hexane)

15 Example 82

4-(5-Chloro-2-thienyl-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]benzamide



20 The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-(N,N-dimethylamino)methyltetraline hydrochloride.

Melting point: 167 - 169°C (crystallization solvent: ethyl acetate-hexane)


25 Example 83

(3'-Chlorobiphenyl-4-yl)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]carboxamide

WO 01/21577

PCT/JP00/06375

214




The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

5 Melting point: 138 - 139° C (crystallization solvent: tetrahydrofuran-isopropyl ether)

Example 84

(2'-Chlorobiphenyl-4-yl)-N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]carboxamide




The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

15 Melting point: 176 - 177° C (crystallization solvent: tetrahydrofuran-hexane)

Example 85

4'-Methyl-N-[6-(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine

WO 01/21577

PCT/JP00/06375

215

obtained in Example 41-2).

¹H-NMR (CDCl₃) δ: 2.25 (6H, s), 2.33 (2H, t, J = 8.1 Hz),
2.41 (3H, s), 2.84 (2H, t, J = 8.1 Hz), 2.98 (2H, s), 6.33
(1H, s), 7.01 (1H, d, J = 7.8 Hz), 7.39 (1H, d, J = 8.4 Hz),
5 7.48 (1H, s), 7.52 (2H, d, J = 7.8 Hz), 7.67 (2H, d, J =
8.1 Hz), 7.84 (1H, s), 7.91 (2H, d, J = 8.1 Hz).

Elemental analysis for C₂₇H₂₈N₂O


Calcd.: C, 81.78; H, 7.12; N, 7.06

Found: C, 81.51; H, 7.22; N, 6.93

10 Melting point: 195 - 196°C (crystallization solvent: ethyl
acetate-diisopropyl ether)

Example 86

4-Cyclohexyl-N-[6-[(N,N-dimethylamino)methyl]-7,8-
15 dihydro-2-naphthalenyl]benzamide



The titled compound was obtained by carrying out the
same operation as in Example 1, using the 6-[(N,N-
dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine
20 obtained in Example 41-2).

¹H-NMR (CDCl₃) δ: 1.20-1.52 (4H, m), 1.71-1.96 (6H, m), 2.25
(6H, s), 2.33 (2H, t, J = 8.1 Hz), 2.50-2.62 (1H, m), 2.84
(2H, t, J = 8.1 Hz), 2.99 (2H, s), 6.33 (1H, s), 7.00 (1H,
d, J = 7.8 Hz), 7.31 (2H, d, J = 8.1 Hz), 7.36 (1H, d, J
25 = 7.8 Hz), 7.46 (1H, brs), 7.75 (1H, s), 7.78 (2H, d, J
= 8.1 Hz).

Melting point: 179 - 181°C (crystallization solvent: ethyl
acetate-diisopropyl ether)


30 Example 87

6-(2,4-Difluorophenyl)-N-[6-[(1-pyrrolidinyl)methyl]-
7,8-dihydro-2-naphthalenyl]nicotinamide

WO 01/21577

PCT/JP00/06375

216



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
 5 obtained in Reference Example 54.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.81 (4H, m), 2.37 (2H, t, $J = 8.1$ Hz), 2.54 (4H, m), 2.86 (2H, t, $J = 8.1$ Hz), 3.18 (2H, s), 6.37 (1H, s), 6.93 (1H, m), 7.04 (2H, m), 7.38 (1H, m), 7.47 (1H, s), 7.77 (1H, s), 7.91 (1H, m), 8.13 (1H, m), 8.24 (1H, m), 9.16 (1H, s).

Elemental analysis for $\text{C}_{27}\text{H}_{26}\text{F}_2\text{N}_3\text{O}$


Calcd.: C, 72.79; H, 5.66; N, 9.43

Found: C, 72.65; H, 5.52; N, 9.73

Melting point: 169 - 170°C (crystallization solvent: ethyl
 15 acetate - diisopropyl ether)

Example 88

4'-Fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 4, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.41 (1H, m), 1.95 (2H, m), 2.25-2.45 (3H, m), 2.36 (6H, s), 2.85-2.94 (3H, m), 7.13 (3H, m), 7.30 (1H, m), 7.46 (1H, s), 7.59 (2H, m), 7.65 (2H, d, $J = 8.1$ Hz), 7.74 (1H, s), 7.93 (2H, d, $J = 8.1$ Hz).

Elemental analysis for $\text{C}_{26}\text{H}_{27}\text{FN}_2\text{O}$

WO 01/21577

PCT/JP00/06375

217

Calcd.: C, 77.58; H, 6.76; N, 6.96

Found: C, 77.72; H, 6.49; N, 6.79

Melting point: 184 - 186°C (crystallization solvent: ethyl acetate - diisopropyl ether)

5

Example 89

(+)-4'-Fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide, and (-)-4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

Optical resolution of 4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide (2.00 g) obtained in Example 88 was conducted by sample-splitting HPLC using a chiral column (Daicel Co., CHIRALCEL OD 500 mmD x 500 mmL; moving phase n-hexane:ethanol = 85:15), to give (+) form (1.00 g; 99.8%ee) and (-) form (0.89 g; >99.9%ee) as powders. The powders obtained were respectively recrystallized using ethyl acetate - diisopropyl ether, to give the (+) form (855 mg) and (-) form (754 mg) of the titled compounds. The optical rotation of both compounds are shown below.

(+)-4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

Optical rotation: $[\alpha]_D = +50.8^\circ$ C=0.494% (methanol)

(-)-4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

Optical rotation: $[\alpha]_D = +51.2^\circ$ C=0.492% (methanol)

30


Example 90

4'-Chloro-N-[3-[(N,N-dimethylamino)methyl]-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide

WO 01/21577

PCT/JP00/06375

218




The titled compound was obtained by carrying out the same operation as in Example 1, using 3-[(N,N-dimethylamino)methyl]-2H-chromen-7-amine obtained in Reference Example 59.

¹H-NMR (CDCl₃) δ: 2.23 (6H, s), 2.97 (2H, s), 4.79 (2H, s), 6.30 (1H, s), 6.96 (1H, d, J = 8.1 Hz), 7.13 (1H, s), 7.20 (1H, d, J = 8.1 Hz), 7.45 (2H, d, J = 8.6 Hz), 7.56 (2H, d, J = 8.6 Hz), 7.66 (2H, d, J = 8.4 Hz), 7.74 (1H, brs), 7.93 (2H, d, J = 8.4 Hz).

Melting point: 199 - 208°C (crystallization solvent: diisopropyl ether)

Example 91

2',4'-Difluoro-N-[3-[(N,N-dimethylamino)methyl]-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 3-[(N,N-dimethylamino)methyl]-2H-chromen-7-amine obtained in Reference Example 59.

¹H-NMR (CDCl₃) δ: 2.23 (6H, s), 2.97 (2H, s), 4.78 (2H, s), 6.29 (1H, s), 6.80-7.10 (2H, m), 6.96 (1H, d, J = 8.1 Hz), 7.13 (1H, s), 7.20 (1H, d, J = 8.1 Hz), 7.40-7.50 (1H, m), 7.62 (2H, d, J = 8.4 Hz), 7.76 (1H, brs), 7.92 (2H, d, J = 8.4 Hz).

Melting point: 200 - 204°C (crystallization solvent: diisopropyl ether)


WO 01/21577

PCT/JP00/06375

219

Example 92

4'-Chloro-N-[6-[(dimethylamino)methyl]-7,8-dihydro-1-naphthalenyl][1,1'-biphenyl]-4-carboxamide



5

The titled compound was obtained in the same manner as in Example 1, using 6-[(dimethylamino)methyl]-7,8-dihydro-1-naphthalenamine obtained in Reference Example 60.

10

$^1\text{H-NMR}$ (CDCl_3) δ : 2.34 (6H, s), 2.36 (2H, t, $J=8.1$ Hz), 2.80 (2H, t, $J=8.1$ Hz), 3.00 (2H, s), 6.38 (1H, s), 6.94 (1H, d, $J=7.8$ Hz), 7.21 (1H, t, $J=7.8$ Hz), 7.45 (2H, d, $J=8.6$ Hz), 7.56 (2H, d, $J=8.6$ Hz), 7.61 (2H, m), 7.68 (2H, d, $J=8.4$ Hz), 7.97 (2H, d, $J=8.4$ Hz).


15

Melting point: 193 - 195°C (crystallization solvent : diisopropyl ether)

Example 93

4'-Chloro-N-[7-[(dimethylamino)methyl]-5,6-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

20



The titled compound was obtained as a white powder by the same method as in Example 1, using 7-[(dimethylamino)methyl]-5,6-dihydro-2-naphthalenamine obtained in Reference Example 61.

25

$^1\text{H-NMR}$ (CDCl_3) δ : 2.25 (6H, s), 2.34 (2H, t, $J=8.1$ Hz), 2.82 (2H, t, $J=8.1$ Hz), 3.00 (2H, s), 6.36 (1H, s), 7.11 (1H, d, $J=7.5$ Hz), 7.34 (1H, d, $J=8.1$ Hz), 7.38 (1H, s), 7.44

WO 01/21577

PCT/JP00/06375

220


(2H, d, J=8.4 Hz), 7.56 (2H, d, J=8.4 Hz), 7.66 (2H, d, J=8.4 Hz), 7.78 (1H, brs), 7.97 (2H, d, J=8.4 Hz).

Melting point: 167 - 169°C (crystallization solvent : diisopropyl ether)

5

Example 94

N-[6-(1-Pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



10 The titled compound was obtained as a white powder in the same manner as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.75-1.90 (4H, m), 2.34 (2H, t, J=8.1 Hz), 2.45-2.60 (4H, m), 2.85 (2H, t, J=8.1 Hz), 3.18 (2H, s), 6.36 (1H, s), 7.02 (1H, d, J=8.1 Hz), 7.27-7.55 (5H, m), 7.63 (2H, d, J=7.3 Hz), 7.70 (2H, d, J=8.4 Hz), 7.82 (1H, s), 7.94 (2H, d, J=8.1 Hz).

15 Elemental analysis for C₂₈H₂₆N₂O


20 Calcd.: C, 82.32; H, 6.91; N, 6.86.

Found: C, 81.99; H, 6.69; N, 6.91.

Melting point: 176 - 177°C (crystallization solvent : diisopropyl ether)

25 Example 95

4'-Fluoro-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



30 The titled compound was obtained in the same manner as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-

WO 01/21577

PCT/JP00/06375

221

dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.75-1.90 (4H, m), 2.35 (2H, t, J=8.2 Hz), 2.45-2.60 (4H, m), 2.84 (2H, t, J=8.2 Hz), 3.18 (2H, s), 6.36 (1H, s), 7.01 (1H, d, J=8.1 Hz), 7.16 (2H, t, J=8.1 Hz), 7.38 (1H, d, J=8.1 Hz), 7.48 (1H, brs), 7.56-7.61 (2H, m), 7.64 (2H, d, J=8.4 Hz), 7.83 (1H, s), 7.93 (2H, d, J=8.4 Hz).

Elemental analysis for C₂₈H₂₇N₂O


Calcd.: C, 78.85; H, 6.38; N, 6.57.

Found: C, 78.75; H, 6.39; N, 6.45.

Melting point: 189 - 192°C (crystallization solvent : diisopropyl ether)

Example 96

N-[6-(1-Pyrrolidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as a white powder in the same manner as in Example 1, using 6-(1-pyrrolidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine obtained in Reference Example 55.

¹H-NMR (CDCl₃) δ: 1.40-1.50 (1H, m), 1.80 (4H, m), 1.80-2.10 (1H, m), 1.80-2.20 (8H, m), 3.30-4.00 (3H, m), 7.29 (1H, d, J=8.4 Hz), 7.25-7.30 (1H, m), 7.30-7.55 (4H, m), 6.43 (2H, d, J=7.0 Hz), 7.70 (2H, t, J=8.4 Hz), 7.75 (1H, s), 7.94 (2H, d, J=8.4 Hz).

Elemental analysis for C₂₈H₃₀N₂O

Calcd.: C, 81.91; H, 7.37; N, 6.82.

Found: C, 81.53; H, 7.25; N, 6.86.

Melting point: 144 - 146°C (crystallization solvent : diisopropyl ether)


WO 01/21577

PCT/JP00/06375

222

Example 97

4'-Fluoro-N-[6-(1-pyrrolidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



5 The titled compound was obtained as a white powder in the same manner as in Example 1, using 6-(1-pyrrolidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine obtained in Reference Example 55.

¹H-NMR (CDCl₃) δ: 1.40-1.50 (1H, m), 1.80 (4H, m), 1.80-2.10 (1H, m), 1.80-2.20 (8H, m), 3.30-4.00 (3H, m), 7.08 (1H, d, J=8.1 Hz), 7.15 (2H, t, J=8.4 Hz), 7.30 (1H, d, J=8.1 Hz), 7.44 (1H, brs), 7.56-7.61 (2H, m), 7.62 (2H, d, J=8.1 Hz), 7.85 (1H, s), 7.92 (2H, d, J=8.1 Hz).

Elemental analysis for C₂₈H₂₉FN₂O


15 Calcd.: C, 78.48; H, 6.82; N, 6.54.

Found: C, 78.18; H, 6.60; N, 6.60.

Melting point: 185 - 189°C (crystallization solvent : diisopropyl ether)

20 Example 98

4'-Chloro-N-[6-(1-pyrrolidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



25 The titled compound was obtained as a white powder in the same manner as in Example 1, using 6-(1-pyrrolidinylmethyl)-5,6,7,8-tetrahydro-2-naphthalenamine obtained in Reference Example 55.

¹H-NMR (CDCl₃) δ: 1.40-1.50 (1H, m), 1.80 (4H, m), 1.80-2.10 (1H, m), 1.80-2.20 (8H, m), 3.30-4.00 (3H, m), 7.08 (1H, d, J=8.1 Hz), 7.31 (1H, d, J=8.4 Hz), 7.43 (2H, d, J=8.7 Hz), 7.45 (1H, s), 7.54 (2H, d, J=8.7 Hz), 7.64 (2H, d, J=8.4

WO 01/21577

PCT/JP00/06375

223

Hz), 7.80 (1H, s), 7.93 (2H, d, J=8.4 Hz).

Elemental analysis for $C_{28}H_{29}ClN_2O$


Calcd.: C, 75.57; H, 6.57; N, 6.30.

Found: C, 75.26; H, 6.68; N, 6.15.

5 Melting point: 206 - 209°C (crystallization solvent :
diisopropyl ether)

Example 99

4-(4-Fluorophenyl)-N-[6-(1-piperidinylmethyl)-7,8-
10 dihydro-2-naphthalenyl]-1-piperidinecarboxamide



6-(1-Pyrrolidinylmethyl)-7,8-dihydro-2-
naphthalenamine obtained in Reference Example 54 (50 mg,
0.22 mmol) and pyridine (35 mg, 0.44 mmol) were dissolved
15 in tetrahydrofuran (3 ml). Phenyl chlorocarbonate (38 mg,
0.24 mol) was added to the solution under ice-cooling, which
was stirred for 10 minutes. The reaction mixture was
concentrated, and dimehtylsulfoxide (5 ml) was added to the
residue. 4-(4-Fluorophenyl)piperidine hydrochloride (57
20 mg, 0.26 mmol) and 4N aqueous sodium hydroxide solution
(0.066 ml, 0.26 mmol) were added to the reaction mixture
at room temperature while stirring, which was stirred for
30 minutes. Ethyl acetate and water were added to the
mixture, and extraction was conducted. The organic layer
25 was washed with water, and concentrated. Diisopropyl
ether was added to the residue. The crystallized product
was collected by filtration, washed with diisopropyl ether,
to give 4-(4-fluorophenyl)-N-[6-(1-piperidinylmethyl)-
7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide (48
30 mg) as a white powder.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.60-1.70 (2H, m), 1.79 (4H, m), 1.80-1.90
(2H, m), 2.33 (2H, t, J=7.8 Hz), 2.51 (4H, m), 2.60-2.70
(1H, m), 2.80 (2H, t, J=7.8 Hz), 2.90-3.10 (2H, m), 3.16

WO 01/21577

PCT/JP00/06375

224


(2H, s), 4.18-4.23 (2H, m), 6.32 (1H, s), 6.32 (1H, s),
6.92-7.09 (4H, m), 7.15-7.20 (3H, m).

Melting point: 182 - 185°C (crystallization solvent :
diisopropyl ether)

5

Example 100

4-(4-Fluorophenyl)-N-[6-(1-pyrrolidinymethyl)-7,8-
dihydro-2-naphthalenyl]-1-piperazinecarboxamide



10

The titled compound was obtained as a white powder in
the same manner as in Example 99, using 6-(1-
pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine
obtained in Reference Example 54 and 4-
fluorophenylpiperazine.

15

¹H-NMR (CDCl₃) δ: 1.79 (4H, m), 2.32 (2H, t, J=7.8 Hz),
2.51 (4H, m), 2.80 (2H, t, J=7.8 Hz), 3.13-3.16 (4H, m), 3.16
(2H, s), 3.63-3.66 (4H, m), 6.30 (1H, s), 6.32 (1H, s),
6.88-7.08 (6H, m), 7.19 (1H, s).

Elemental analysis for C₂₆H₃₁FN₄O

20

Calcd.: C, 71.86; H, 7.19; N, 12.89.


Found: C, 71.68; H, 7.35; N, 12.65.

Melting point: 179 - 181°C (crystallization solvent :
diisopropyl ether)

25

Example 101

N-(4-Bromophenyl)-6-[(dimethylamino)methyl]-7,8-
dihydro-2-naphthalenecarboxamide



1) 6-Cyano-1-tetralone (1.30 g, 7.59 mmol)

30

synthesized by a known method by documents (synthetic
communications, 23(21), 2965 (1993)) was dissolved in a

mixed solution of concentrated hydrochloric acid (10 ml) and acetic acid (20 ml), which was stirred at 120°C for 16 hours. The reaction mixture was concentrated. Ethyl acetate and water were added to the residue, and extraction
5 was conducted. The organic layer was washed with water, and concentrated. The residue was washed with ethyl acetate - n-hexane (1:1), to give 5-oxo-5,6,7,8-tetrahydro-2-naphthalenecarboxylic acid (1.10 g) as a white powder.

10 ¹H-NMR (CDCl₃) δ: 2.15-2.23 (2H, m), 2.70-2.75 (2H, m), 3.04-3.07 (2H, m), 8.01-8.03 (1H, m), 8.03 (1H, s), 8.13 (1H, d, J=8.7 Hz).

2) N-(4-Bromophenyl)-5-oxo-5,6,7,8-tetrahydro-2-naphthalenecarboxamide (1.21 g) was obtained as a white
15 powder in the same manner as in Example 1, using 5-oxo-5,6,7,8-tetrahydro-2-naphthalenecarboxylic acid (1.00 g, 5.26 mmol) obtained in 1) and 4-bromoaniline (0.90 g, 5.26 mmol).

20 ¹H-NMR (CDCl₃) δ: 2.14-2.23 (2H, m), 2.69-2.73 (2H, m), 3.03-3.07 (2H, m), 7.48-7.58 (4H, m), 7.71 (1H, d, J=8.1 Hz), 7.79 (1H, s), 7.86 (1H, s), 8.12 (1H, d, J=8.1 Hz).

3) N-(4-Bromophenyl)-5-oxo-5,6,7,8-tetrahydro-2-naphthalenecarboxamide (1.10 g, 3.19 mmol) obtained in 2) was dissolved in dimethylformamide diethylacetal (30 ml),
25 which was refluxed with heating for 4 hours. The crystallized product was collected by filtration, washed with ethyl acetate, to give N-(4-bromophenyl)-6-[(dimethylamino)methylidene]-5-oxo-5,6,7,8-tetrahydro-2-naphthalenecarboxamide (1.21 g) as a yellow powder.

30 ¹H-NMR (CDCl₃) δ: 2.80-2.87 (4H, m), 3.07 (6H, m), 7.46-7.72 (7H, m), 7.91 (1H, d, J=8.4 Hz), 8.53 (1H, s).

4) Sodium triacetoxyhydroborate (398 mg, 1.87 mmol) was dissolved in a mixed solution of acetic acid (40 ml) and tetrahydrofuran (10 ml) under ice-cooling. N-(4-
35 Bromophenyl)-6-[(dimethylamino)methylidene]-5-oxo-5,6,7,8-tetrahydro-2-naphthalenecarboxamide (500 mg,

1.25 mmol) obtained in 3) was added to the solution, which was stirred for 1 hour. The reaction mixture was concentrated under reduced pressure at room temperature.

2-Propanol (50 ml) was added to the residue, and sodium borohydride (142 mg, 3.75 mmol) was further added under ice-cooling. After stirring for 2 hours, the reaction mixture was concentrated. Sodium hydrogencarbonate solution and ethyl acetate was added to the residue for liquid separation. The organic layer was concentrated. The residue was dissolved in a mixed solution of acetic acid (20 ml) and concentrated hydrochloric acid (20 ml), which was stirred at 70°C for 5 hours. The reaction mixture was concentrated. 4N aqueous sodium hydroxide solution and ethyl acetate were added to the residue, and extraction was conducted. The organic layer was washed with water, and concentrated. The residue was purified by alumina column chromatography (development solvent: ethyl acetate), and the eluent was washed with diisopropyl ether, to give the titled compound (234 mg) as a white powder.

¹H-NMR (CDCl₃) δ: 2.26 (6H, s), 2.38 (2H, t, J=8.1 Hz), 2.89 (2H, t, J=8.1 Hz), 3.02 (2H, s), 6.42 (1H, s), 7.10 (1H, d, J=8.6 Hz), 7.47 (2H, d, J=8.9 Hz), 7.55 (2H, d J=8.9 Hz), 7.61 (1H, s), 7.62 (1H, d, J=6.7 Hz), 7.76 (1H, s).

Elemental analysis for C₂₀H₂₁BrN₂O


Calcd.: C, 62.35; H, 5.49; N, 7.27.

Found: C, 61.98; H, 5.43; N, 7.07.

Melting point: 175 - 179°C (crystallization solvent : diisopropyl ether)

Example 102

6-[(Dimethylamino)methyl]-N-(4'-fluoro[1,1'-biphenyl]-4-yl)-7,8-dihydro-2-naphthalenecarboxamide



WO 01/21577

PCT/JP00/06375

227


The titled compound was obtained as a white powder, by the same method as in Example 16, using N-(4-bromophenyl)-6-[(dimethylamino)methyl]-7,8-dihydro-2-naphthalenecarboxamide (170 mg, 0.44 mmol) obtained in
5 Example 101 and 4-fluorophenylboric acid (74 mg, 0.53 mmol).

¹H-NMR (CDCl₃) δ: 2.27 (6H, s), 2.39 (2H, t, J=8.4 Hz), 2.91 (2H, t, J=8.4 Hz), 3.02 (2H, s), 6.43 (1H, s), 7.09-7.16 (3H, m), 7.52-7.73 (8H, m), 7.81 (1H, s).

10 Melting point: 200 - 204°C (crystallization solvent : diisopropyl ether)

Example 103

2',4'-Difluoro-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-
15 2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as a white powder by the same method as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
20 obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.75-1.90 (4H, m), 2.36 (2H, t, J=8.1 Hz), 2.45-2.60 (4H, m), 2.85 (2H, t, J=8.1 Hz), 3.18 (2H, s), 6.36 (1H, s), 6.92-7.03 (3H, m), 7.36-7.45 (2H, m), 7.48 (1H, s), 7.62 (2H, d, J=8.4 Hz), 7.78 (1H, s), 7.94 (2H, d, J=8.4 Hz).
25

Elemental analysis for C₂₆H₂₆F₂N₂O

Calcd.: C, 75.66; H, 5.90; N, 6.30.

Found: C, 75.36; H, 5.92; N, 6.10.

Melting point: 165 - 167°C (crystallization solvent :
30 diisopropyl ether)

Example 104


N-[3-[(Dimethylamino)methyl]-2,3-dihydro-1,4-

WO 01/21577

PCT/JP00/06375

228

benzodioxin-6-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as a white powder by the same method as in Example 1, using N,N-dimethyl-N-[(7-amino-2,3-dihydro-1,4-benzodioxin-2-yl)methyl]amine obtained in Reference Example 62.

¹H-NMR (CDCl₃) δ: 2.33 (6H, s), 2.48-2.66 (2H, m), 3.93-3.99 (1H, m), 4.27-4.31 (2H, m), 6.86 (1H, d, J=8.6 Hz), 7.03-7.07 (1H, m), 7.31-7.32 (1H, m), 7.37-7.49 (3H, m), 7.62 (2H, d, J=7.0 Hz), 7.68 (2H, d, J=8.4 Hz), 7.76 (1H, s), 7.91 (2H, d, J=8.4 Hz).

Elemental analysis for C₂₄H₂₄N₂O₃


Calcd.: C, 74.21; H, 6.23; N, 7.21.

Found: C, 74.17; H, 6.23; N, 7.01.

Melting point: 124 - 126°C (crystallization solvent : diisopropyl ether)

Example 105

4'-Chloro-N-[3-[(dimethylamino)methyl]-2,3-dihydro-1,4-benzodioxin-6-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as a white powder by the same method as in Example 1, using N,N-dimethyl-N-[(7-amino-2,3-dihydro-1,4-benzodioxin-2-yl)methyl]amine obtained in Reference Example 62.

¹H-NMR (CDCl₃) δ: 2.33 (6H, s), 2.50-2.67 (2H, m), 3.94-4.01 (1H, m), 4.28-4.31 (2H, m), 6.86 (1H, d, J=8.7 Hz), 7.03-7.06 (1H, m), 7.31 (1H, m), 7.44 (2H, d, J=8.4 Hz), 7.55 (2H, d, J=8.4 Hz), 7.65 (2H, d, J=8.1 Hz), 7.67 (1H, s), 7.91 (2H, d, J=8.1 Hz).

WO 01/21577


PCT/JP00/06375

229

Melting point: 158 - 159°C (crystallization solvent :
diisopropyl ether)

Example 106

5 4'-Chloro-N-[2-[(dimethylamino)methyl]-2,3-dihydro-1,4-benzodioxin-6-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as a white powder by
the same method as in Example 1, using N,N-dimethyl-N-
10 [(6-amino-2,3-dihydro-1,4-benzodioxin-2-yl)methyl]amine
obtained in Reference Example 63.

¹H-NMR (CDCl₃) δ: 2.34 (6H, s), 2.46-2.67 (2H, m), 3.94-4.01
(1H, m), 4.28-4.34 (2H, m), 6.91 (1H, d, J=8.6 Hz),
7.02-7.05 (1H, m), 7.30 (1H, m), 7.44 (2H, d, J=8.4 Hz),
15 7.55 (2H, d, J=8.4 Hz), 7.66 (2H, d, J=8.1 Hz), 7.70 (1H,
s), 7.92 (2H, d, J=8.1 Hz).

Elemental analysis for C₂₄H₂₃ClN₂O₃


Calcd.: C, 68.16; H, 5.48; N, 6.62.

Found: C, 68.09; H, 5.29; N, 6.57.

20 Melting point: 215 - 217°C (crystallization solvent :
diisopropyl ether)

Example 107

25 2',4'-Difluoro-N-[2-[(dimethylamino)methyl]-2,3-dihydro-1,4-benzodioxin-6-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as a white powder by
the same method as in Example 1, using N,N-dimethyl-N-
30 [(6-amino-2,3-dihydro-1,4-benzodioxin-2-yl)methyl]amine

WO 01/21577

PCT/JP00/06375

230

obtained in Reference Example 63.

¹H-NMR (CDCl₃) δ: 2.34 (6H, s), 2.50-2.63 (2H, m), 3.94-4.01 (1H, m), 4.28-4.34 (2H, m), 6.91 (1H, d, J=8.6 Hz), 6.91-7.03 (3H, m), 7.30 (1H, m), 7.40-7.50 (1H, m), 7.61 (2H, d, J=8.1 Hz), 7.69 (1H, s), 7.92 (2H, d, J=8.1 Hz).

Elemental analysis for C₂₄H₂₂F₂N₂O₃


Calcd.: C, 67.91; H, 5.22; N, 6.60.

Found: C, 67.75; H, 5.09; N, 6.48.

Melting point: 209 - 210°C (crystallization solvent : diisopropyl ether)

Example 108

6-(4-Chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-2,3-dihydro-1,4-benzodioxin-6-yl]nicotinamide



The titled compound was obtained as a white powder by the same method as in Example 1, using 1-[(6-amino-2,3-dihydro-1,4-benzodioxin-2-yl)methyl]pyrrolidine obtained in Reference Example 64.

¹H-NMR (CDCl₃) δ: 1.81 (4H, m), 2.50-2.63 (4H, m), 2.75-2.77 (2H, m), 3.90-4.10 (1H, m), 4.30-4.36 (2H, m), 6.91 (1H, d, J=8.6 Hz), 7.00-7.10 (1H, m), 7.26 (1H, m), 7.48 (2H, d, J=8.6 Hz), 7.72 (1H, s), 7.81 (1H, d, J=7.8 Hz), 8.01 (2H, d, J=8.6 Hz), 8.20-8.25 (1H, m), 9.10 (1H, s).

Elemental analysis for C₂₅H₂₄ClN₃O₃

Calcd.: C, 66.74; H, 5.38; N, 9.34.

Found: C, 66.66; H, 5.46; N, 9.11.

Melting point: 218 - 220°C (crystallization solvent : diisopropyl ether)


Example 109

N-[3-[(Dimethylamino)methyl]-2H-chromen-7-yl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide

WO 01/21577

PCT/JP00/06375

231



The titled compound was obtained by carrying out the same operation as in Example 1, using 3-[(N,N-dimethylamino)methyl]-2H-chromen-7-amine obtained in Reference Example 59.

¹H-NMR (CDCl₃) δ: 2.23 (6H, s), 2.97 (2H, s), 4.79 (2H, s), 6.30 (1H, s), 6.96 (1H, d, J=8.1 Hz), 7.13-7.22 (4H, m), 7.56-7.61 (2H, m), 7.65 (2H, d, J=8.4 Hz), 7.78 (1H, s), 7.92 (2H, d, J=8.4 Hz).

Elemental analysis for C₂₅H₂₃N₂O₂

Calcd.: C, 74.61; H, 5.76; N, 6.96.


Found: C, 74.35; H, 5.68; N, 6.74.

Melting point: 192 - 195°C (crystallization solvent : diisopropyl ether)

15

Example 110

4'-Chloro-N-[3-[(dimethylamino)methyl]-3,4-dihydro-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using N-[(7-amino-3,4-dihydro-2H-chromen-3-yl)methyl]-N,N-dimethylamine obtained in Reference Example 65.

¹H-NMR (CDCl₃) δ: 2.26 (6H, s), 2.27 (3H, m), 2.47-2.51 (1H, m), 2.83-2.89 (1H, m), 3.82-3.86 (1H, m), 4.28-4.32 (1H, m), 7.04 (1H, d, J=8.1 Hz), 7.12-7.18 (2H, m), 7.44 (2H, d, J=8.4 Hz), 7.56 (2H, d, J=8.4 Hz), 7.67 (2H, d, J=8.4 Hz), 7.71 (1H, s), 7.93 (2H, d, J=8.4 Hz).


30 Example 111

WO 01/21577

PCT/JP00/06375

232

4'-Chloro-N-[6-[(dimethylamino)methyl]-5-methyl-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-

[(dimethylamino)methyl]-5-methyl-7,8-dihydro-2-naphthalenamine obtained in Reference Example 66.

¹H-NMR (CDCl₃) δ: 2.09 (3H, s), 2.27 (6H, s), 2.31-2.37 (2H, m), 2.74-2.79 (2H, m), 3.08 (2H, s), 7.27-7.30 (1H, m), 7.44-7.48 (4H, m), 7.56 (2H, d, J=8.6 Hz), 7.67 (2H, d, J=8.4 Hz), 7.79 (1H, s), 7.95 (2H, d, J=8.4 Hz).

Elemental analysis for C₂₇H₂₇ClN₂O


Calcd.: C, 75.25; H, 6.31; N, 6.50.

Found: C, 74.86; H, 6.20; N, 6.42.

Melting point: 199 - 204°C (crystallization solvent : diisopropyl ether)

Example 112

4'-Chloro-N-[6-[(dimethylamino)methyl]-5-ethyl-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-

[(dimethylamino)methyl]-5-ethyl-7,8-dihydro-2-naphthalenamine obtained in Reference Example 67.

¹H-NMR (CDCl₃) δ: 1.09 (3H, t, J=7.5 Hz), 2.27 (6H, s), 2.31-2.37 (2H, m), 2.60-2.63 (2H, m), 2.71-2.76 (2H, m), 3.08 (2H, s), 7.31 (1H, d, J=9.2 Hz), 7.43-7.49 (4H, m),

WO 01/21577

PCT/JP00/06375

233

7.56 (2H, d, J=8.7 Hz), 7.67 (2H, d, J=8.6 Hz), 7.80 (1H, s), 7.94 (2H, d, J=8.6 Hz).

Elemental analysis for $C_{28}H_{29}ClN_2O$


Calcd.: C, 75.57; H, 6.57; N, 6.30.

5 Found: C, 75.41; H, 6.34; N, 6.23.

Melting point: 201 - 204°C (crystallization solvent : diisopropyl ether)

Example 113

10 4'-Chloro-N-[6-[(dimethylamino)methyl]-5-isobutyl-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-

15 [(dimethylamino)methyl]-5-isobutyl-7,8-dihydro-2-naphthalenamine obtained in Reference Example 68.

1H -NMR ($CDCl_3$) δ : 0.90 (6H, d, J=6.4 Hz), 1.73-1.78 (1H, m), 2.23 (6H, s), 2.34 (2H, m), 2.50 (2H, d, J=7.3 Hz), 2.74 (2H, m), 3.13 (2H, s), 7.26-7.30 (1H, m), 7.45-7.48 (4H, m), 7.56 (2H, d, J=8.7 Hz), 7.67 (2H, d, J=8.4 Hz), 7.79 (1H, s), 7.94 (2H, d, J=8.4 Hz).

20 Elemental analysis for $C_{30}H_{33}ClN_2O$

Calcd.: C, 76.17; H, 7.03; N, 5.92.

Found: C, 75.91; H, 7.19; N, 5.72.

25 Melting point: 159 - 162°C (crystallization solvent : diisopropyl ether)


Example 114

30 4'-Chloro-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

WO 01/21577

PCT/JP00/06375

234



The titled compound was obtained by carrying out the same operation as in Example 1, using 5-methyl-6- (1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
5 obtained in Reference Example 69.


¹H-NMR (CDCl₃) δ: 1.79 (4H, m), 2.11 (3H, s), 2.30-2.40 (2H, m), 2.54 (4H, m), 2.74-2.79 (2H, m), 3.28 (2H, s), 7.26-7.30 (1H, m), 7.45-7.48 (4H, m), 7.56 (2H, d, J=8.6 Hz), 7.67 (2H, d, J=8.4 Hz), 7.81 (1H, s), 7.95 (2H, d, J=8.4 Hz).

10

Melting point: 190 - 192°C (crystallization solvent : diisopropyl ether)

Example 115

15 N-[5-Methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 5-methyl-6- (1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
20 obtained in Reference Example 69.

¹H-NMR (CDCl₃) δ: 1.78 (4H, m), 2.10 (3H, s), 2.35-2.40 (2H, m), 2.53 (4H, m), 2.70-2.78 (2H, m), 3.28 (2H, s), 7.26-7.28 (1H, m), 7.40-7.50 (5H, m), 7.62 (2H, d, J=7.0 Hz), 7.70 (2H, d, J=8.4 Hz), 7.87 (1H, s), 7.94 (2H, d, J=8.4 Hz).

25

Melting point: 169 - 170°C (crystallization solvent : diisopropyl ether)


WO 01/21577

PCT/JP00/06375

235

Example 116

6-(4-Methoxyphenyl)-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 69.

10 $^1\text{H-NMR}$ (CDCl_3) δ : 1.78 (4H, m), 2.09 (3H, s), 2.35-2.40 (2H, m), 2.53 (4H, m), 2.70-2.77 (2H, m), 3.27 (2H, s), 3.88 (3H, s), 7.01 (2H, d, $J=8.9$ Hz), 7.26 (1H, d, $J=8.9$ Hz), 7.45-7.47 (2H, m), 7.75 (1H, d, $J=8.4$ Hz), 7.95 (1H, s), 8.01 (2H, d, $J=8.9$ Hz), 8.18-8.21 (1H, m), 9.09 (1H, m).

15 Elemental analysis for $\text{C}_{29}\text{H}_{31}\text{N}_3\text{O}_2$

Calcd.: C, 76.79; H, 6.89; N, 9.26.


Found: C, 76.46; H, 6.64; N, 9.09.

Melting point: 165 - 167°C (crystallization solvent : diisopropyl ether)

20

Example 117

4'-Chloro-N-[5-cyano-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



25 The titled compound was obtained as a colorless powder by carrying out the same operation as in Example 1, using 6-amino-2-(1-pyrrolidinylmethyl)-3,4-dihydro-1-naphthalenecarbonitrile obtained in Reference Example 70 and 4'-chloro[1,1'-biphenyl]-4-carboxylic acid.

WO 01/21577

PCT/JP00/06375

236


^1H NMR ($\text{DMSO}-d_6$) δ : 1.73 (4H, m), 2.50 (4H, m), 2.56 (2H, m), 2.82 (2H, m), 3.49 (2H, s), 7.32 (1H, d, $J = 9.0$ Hz), 7.57 (2H, d, $J = 8.4$ Hz), 7.56-7.87 (6H, m), 8.07 (2H, d, $J = 8.4$ Hz), 10.40 (1H, s).

5 FABMS(pos) 468.2 $[\text{M}+\text{H}]^+$

Melting point: 191 - 192°C (crystallization solvent : diisopropyl ether)

Example 118

10 N-[5-Cyano-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by as a colorless powder carrying out the same operation as in Example 1, using 6-amino-2-(1-pyrrolidinylmethyl)-3,4-dihydro-1-naphthalenecarbonitrile obtained in Reference Example 70 and [1,1'-biphenyl]-4-carboxylic acid.

15 ^1H NMR ($\text{DMSO}-d_6$) δ : 1.81 (4H, m), 2.62 (6H, m), 2.88 (2H, m), 3.56 (2H, s), 7.41 (2H, m), 7.46 (3H, m), 7.64 (2H, d, $J = 6.9$ Hz), 7.73 (3H, m), 7.88 (1H, s), 7.95 (2H, d, $J = 8.1$ Hz).


20 FABMS(pos) 434.2 $[\text{M}+\text{H}]^+$

Melting point: 168 - 170°C (crystallization solvent : diisopropyl ether)

25

Example 119

3-Bromo-N-[6-[(dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl]benzamide




30 The titled compound was obtained by carrying out the same operation as in Example 1, using 6-amino-2-[(N,N-

dimethylamino)methyl]tetralin and 3-bromobenzoic acid.

¹H NMR (DMSO-d₆) δ: 1.31 (1H, m), 1.89 (2H, m), 2.17 (6H, s), 2.17-2.35 (3H, m), 2.77 (3H, m), 7.04 (1H, d, J=8.4 Hz), 7.49 (3H, m), 7.79 (1H, d, J=8.1 Hz), 7.94 (1H, d, J=7.8 Hz), 8.13 (1H, s), 10.20 (1H, s).

Example 120

N-[6-[(Dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-3-carboxamide



10

The titled compound was obtained by carrying out the same operation as in Example 16, using 3-bromo-N-[6-[(dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl]benzamide obtained in Example 119 and phenylboronic acid.

15

¹H NMR (DMSO-d₆) δ: 1.43 (1H, m), 2.02 (1H, m), 2.21 (1H, m), 2.42 (1H, m), 2.81 (6H, s), 2.88 (3H, m), 3.09 (2H, m), 7.06 (1H, m), 7.42-7.65 (6H, m), 7.78-7.95 (4H, m), 8.22 (1H, s), 10.27 (1H, s).


20

FABMS(pos) 385.2 [M+H]⁺

Melting point: 145 - 148°C (crystallization solvent : ethyl acetate-diisopropyl ether)

Example 121

N-[6-[(Dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl]-2',4'-difluoro[1,1'-biphenyl]-4-carboxamide



30

The titled compound was obtained by carrying out the same operation as in Example 1, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin and 2',4'-difluoro[1,1'-

WO 01/21577

PCT/JP00/06375

238


biphenyl]-4-carboxylic acid.

¹H NMR (CDCl₃) δ: 1.41 (1H, m), 1.94 (2H, m), 2.25 (6H, s),
2.23-2.30 (3H, m), 2.86 (3H, m), 6.96 (2H, m), 7.09 (1H,
d, J=8.1 Hz), 7.30 (1H, m), 7.43 (2H, m), 7.61 (2H, m), 7.76
5 (1H, s), 7.93 (2H, m).

Melting point: 162 - 163°C (crystallization solvent :
ethyl acetate-diisopropyl ether)

Example 122

10 N-[6-[(Dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl]-1H-indole-2-carboxamide



The titled compound was obtained by carrying out the
same operation as in Example 1, using 6-amino-2-[(N,N-
15 dimethylamino)methyl]tetralin and 1H-indol-2-carboxylic
acid.


¹H NMR (DMSO-d₆) δ: 1.32 (1H, m), 1.91 (2H, m), 2.16 (6H,
s), 2.16-2.35 (3H, m), 2.78 (3H, m), 7.06 (2H, m), 7.21 (1H,
m), 7.44 (4H, m), 7.66 (1H, d, J=8.1 Hz), 10.05 (1H, s),
20 11.68 (1H, s).

FABMS(pos) 348.2 [M+H]⁺

Melting point: 190 - 192°C (crystallization solvent :
ethyl acetate - diisopropyl ether)

25 Example 123

N-[6-[(Dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl] [1,1'-biphenyl]-4-carboxamide




A tetrahydrofuran solution (0.146ml, 0.293mmol) of
30 N-(6-oxo-5,6,7,8-tetrahydro-2-naphthalenyl)[1,1'-

biphenyl]-4-carboxamide (10 mg, 0.029 mmol) obtained in Reference Example 72 and 2N dimethylamine was added to acetic acid-tetrahydrofuran (1:1) solution (0.5ml), which was stirred at 50°C for 15 minutes. After the reaction mixture was cooled at room temperature, sodium triacetoxhydroborate (31 mg, 0.146 mmol) was added, which was stirred at 50°C for 2 hours. 1N Hydrochloric acid was added to the reaction mixture, which was washed with ethyl acetate. Sodium carbonate was added to the water layer to make it alkaline, then extraction was conducted using ethyl acetate. The extract was washed with saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. The resulting residue was purified by alumina B column chromatography (development solvent; ethyl acetate), to give the titled compound (1.6mg).
¹H NMR (CDCl₃) δ: 1.68 (1H, m), 2.27 (1H, m), 2.40 (6H, s), 2.78 (5H, m), 7.11 (1H, d, J=8.1 Hz), 7.32-7.50 (5H, m), 7.62 (2H, m), 7.72 (2H, d, J=8.4 Hz), 7.78 (1H, br), 7.94 (2H, d, J=8.4 Hz).
 FABMS(pos) 371.2 [M+H]⁺

Example 124

N-[4-[(E)-2-(4,5-Dihydro-1H-imidazol-2-yl)ethenyl]phenyl]phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride




10.1 N Hydrogen chloride-ethanol solution (30 ml) was added to an ethanol suspension of N-[4-[(E)-2-cyanoethenyl]phenyl][1,1'-biphenyl]-4-carboxamide (250 mg, 0.771 mmol) obtained in Reference Example under room temperature, which was stirred for 16 hours. After the

- solvent was distilled out under reduced pressure, ethanol was again added to the residue, and then ethylenediamine (0.155 ml, 2.31 mmol) was added at room temperature, which was stirred for 16 hours. Sodium hydrogencarbonate solution was added to the reaction mixture, and the precipitated crude product was washed with water and chloroform. This product was dissolved in methanol. 1 N Hydrochloric acid (4 ml) was added to the solution, and the solvent was distilled out under reduced pressure.
- Small amount of water was added to the resulting residue, to give the titled compound (124 mg) as a colorless powder. ^1H NMR (DMSO- d_6 , free base) δ : 3.33 (4H, m), 6.61 (1H, d, $J = 16.8$ Hz), 7.15 (1H, d, $J = 16.8$ Hz), 7.52 (5H, m), 7.83 (6H, m), 8.07 (2H, d, $J = 8.4$ Hz).
- Elemental analysis for $\text{C}_{24}\text{H}_{21}\text{N}_3\text{O} \cdot \text{HCl} \cdot 1.5\text{H}_2\text{O}$
Calcd.: C, 66.89; H, 5.85; N, 9.75.
Found: C, 67.16; H, 6.10; N, 10.03.

Example 125

- N-[4-[2-(4,5-Dihydro-1H-imidazol-2-yl)ethenyl]phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride



- 10% Palladium - carbon (200 mg) was added to a methanol suspension of N-[4-[(E)-2-(4,5-dihydro-1H-imidazol-2-yl)ethenyl]phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride (80 mg, 0.198 mmol) obtained in Example 124, which was stirred under hydrogen atmosphere at 60°C for 2 hours. After a catalyst was filtered off, the solvent was distilled out under reduced pressure. Diethyl ether was added to the resulting residue, to give the titled compound (52 mg) as a colorless powder.

WO 01/21577

PCT/JP00/06375

241


^1H NMR (DMSO- d_6) δ : 2.73-2.97 (4H, m), 3.37 (4H, s), 7.24 (2H, d, $J = 8.4$ Hz), 7.46 (3H, m), 7.76 (6H, m), 8.08 (2H, d, $J = 8.4$ Hz).

FABMS(pos) 370[M+H] $^+$

5

Example 126

4-Chloro-N-[2-[[6-[(dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl]amino]-2-oxoethyl]benzamide



10 The titled compound was obtained by carrying out the same operation as in Example 1, using 6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine obtained in Example 41-2) and 4-chlorobenzoyl glycine.


15 ^1H NMR (DMSO- d_6) δ : 2.18 (6H, s), 2.21 (2H, m), 2.71 (2H, m), 2.91 (2H, s), 4.05 (2H, d, $J=5.6$ Hz), 6.30 (1H, s), 6.98 (1H, d, $J=8.1$ Hz), 7.36 (2H, m), 7.58 (2H, d, $J=8.4$ Hz), 7.92 (2H, d, $J=8.4$ Hz), 8.94 (1H, t, $J=5.6$ Hz), 10.00 (1H, s).

FABMS(pos) 398 [M+H] $^+$

20 Melting point: 168 - 171°C (crystallization solvent : diisopropyl ether)

Example 127

25 4'-Chloro-N-[4-(3-piperidinylcarbonyl)phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride



1) tert-Butyl 3-[4-[[[4'-chloro[1,1'-biphenyl]-4-yl]carbonyl]amino]benzoyl]-1-piperidinecarboxylate was obtained by carrying out the same operation as in Example 30 1, using tert-butyl 3-(4-aminobenzoyl)-1-piperidinecarboxylate obtained in Reference Example 77 and

WO 01/21577

PCT/JP00/06375

242


4'-chloro[1,1'-biphenyl]-4-carboxylic acid.

FABMS(pos) 519.2 [M+H]⁺

- 2) 4N Hydrogen chloride-ethyl acetate (1 ml) was added to tert-butyl 3-[4-[[[(4'-chloro[1,1'-biphenyl]-4-yl)carbonyl]amino]benzoyl]-1-piperidinecarboxylate (100 mg, 0.193 mmol) obtained in 1). One hour later, the solvent was distilled out under reduced pressure. Diisopropyl ether was added to the residue, to give the titled compound (73.3 mg) as a colorless powder.
- ¹H NMR (DMSO-d₆) δ : 1.56 (1H, m), 1.82 (2H, m), 2.02 (1H, m), 2.89 (1H, m), 3.05 (1H, m), 3.33 (2H, m), 3.90 (1H, m), 7.58 (2H, d, J=8.1Hz), 7.81 (2H, d, J=8.1Hz), 7.88 (2H, d, J=8.1Hz), 8.03 (4H, m), 8.11 (2H, d, J=8.1Hz), 9.04 (2H, br), 10.73 (1H, s).
- FABMS(pos) 419.2 [M+H]⁺
- Melting point: 222 - 225°C (decomposition)

Example 128

- 4'-Chloro-N-[4-[hydroxy(3-piperidinyl)methyl]phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride



- 4N Hydrogen chloride-ethyl acetate (1 ml) was added to tert-butyl 3-[[4-[[[(4'-chloro[1,1'-biphenyl]-4-yl)carbonyl]amino]phenyl](hydroxy)methyl]-1-piperidinecarboxylate (100 mg, 0.192 mmol) obtained in Reference Example 78. One hour later, the solvent was distilled out under reduced pressure. Diisopropyl ether was added to the residue, to give the titled compound (79.8 mg) as a colorless powder.
- FABMSMS(pos) 421.2 [M+H]⁺
- Melting point: 195°C (decomposition)


WO 01/21577

PCT/JP00/06375

243

Example 129

[4-[[[4'-Chloro[1,1'-biphenyl]-4-yl)carbonyl]amino]phenyl](3-piperidiny)methyl acetate



Concentrated sulfuric acid (0.0562 ml) was added to an acetic acid solution (3.5 ml) of tert-butyl 3-[[4-[[[4'-chloro[1,1'-biphenyl]-4-yl)carbonyl]amino]phenyl](hydroxy)methyl]-1-piperidinecarboxylate (366 mg, 0.702 mmol) obtained in Example 128, which was stirred under room temperature for 16 hours. Ethyl acetate was added to the reaction mixture, which was washed with potassium hydrogencarbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. The resulting oily substance was purified by alumina B column chromatography (development solvent; ethyl acetate: methanol = 3:1), and powdered with diisopropyl ether, to give the titled compound (210 mg).


10
15
20

FABMS(pos) 403.2 [M+H]⁺

Melting point: 200 - 203°C.

Example 130

25 N-[4-(3-Piperidinylmethyl)phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride



4N Hydrogen chloride-ethyl acetate (2 ml) was added to tert-butyl 3-[4-[[[1,1'-biiphenyl]-4-

WO 01/21577

PCT/JP00/06375

244


ylcarbonyl)amino]benzyl]-1-piperidinecarboxylate (100 mg, 0.212 mmol) obtained in Reference Example 80. Two hours later, the solvent was distilled out under reduced pressure. Diisopropyl ether was added to the residue for powdering, to give the titled compound (79 mg).

FABMS(pos) 371.3 [M+H]⁺

Melting point: 218 - 220°C (decomposition)

Example 131

4'-Fluoro-N-[4-(3-piperidinylmethyl)phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride




4N Hydrogen chloride-ethyl acetate (3 ml) was added to tert-butyl 3-[4-[[[(4'-fluoro[1,1'-biphenyl]-4-yl)carbonyl]amino]benzyl]-1-piperidinecarboxylate (150 mg, 0.307 mmol) obtained in Reference Example 81. Two hours later, the solvent was distilled out under reduced pressure. Diisopropyl ether was added to the residue, to give the titled compound (115 mg) as a colorless powder.

FABMS(pos) 389.3 [M+H]⁺

Melting point: 205°C (decomposition)

Example 132

4'-Chloro-N-[4-(3-piperidinylmethyl)phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride



4N Hydrogen chloride-ethyl acetate (3 ml) was added to tert-butyl 3-[4-[[[(4'-chloro[1,1'-biphenyl]-4-yl)carbonyl]amino]benzyl]-1-piperidinecarboxylate (150 mg, 0.297 mmol) obtained in Reference Example 82. Two hours

WO 01/21577

PCT/JP00/06375

245


later, the solvent was distilled out under reduced pressure. Diisopropyl ether was added to the residue, to give the titled compound (73.3 mg) as a colorless powder.

FABMS(pos) 405.2 [M+H]⁺

5 Melting point: 200°C (decomposition)

Example 133

N-[7-[(Dimethylamino)methyl]-5,6-dihydro-3-quinolinyl][1,1'-biphenyl]-4-carboxamide



10

The titled compound was obtained by carrying out the same operation as in Example 1, using N-[(3-amino-5,6-dihydro-7-quinolinyl)methyl]-N,N-dimethylamine obtained in Reference Example 86 and [1,1'-biphenyl]-4-carboxylic acid.

15

¹H NMR (DMSO-d₆) δ : 2.16 (6H, s), 2.29 (2H, t, J=8.1 Hz), 2.84 (2H, t, J=8.1 Hz), 2.98 (2H, s), 6.40 (1H, s), 7.42 (1H, m), 7.51 (2H, m), 7.76 (2H, d, J=7.2 Hz), 7.84 (2H, d, J=8.1 Hz), 7.97 (1H, s), 8.06 (2H, d, J=8.4 Hz), 8.65 (1H, s), 10.39 (1H, s).


20

FABMS(pos) 384.2 [M+H]⁺

Melting point: 202 - 203°C.

Example 134

25 4'-Chloro-N-[7-[(dimethylamino)methyl]-5,6-dihydro-3-quinolinyl][1,1'-biphenyl]-4-carboxamide



30

The titled compound was obtained by carrying out the same operation as in Example 1, using N-[(3-amino-5,6-dihydro-7-quinolinyl)methyl]-N,N-dimethylamine obtained

WO 01/21577

PCT/JP00/06375

246

in Reference Example 86 and 4'-chloro[1,1'-biphenyl]-4-carboxylic acid.

¹H NMR (DMSO-d₆) δ: 2.17 (6H, s), 2.31 (2H, t, J=8.1 Hz), 2.85 (2H, t, J=8.1 Hz), 2.99 (2H, s), 6.41 (1H, s), 7.57 (2H, d, J=8.4 Hz), 7.81 (2H, d, J=8.4 Hz), 7.86 (2H, d, J=8.4 Hz), 7.98 (1H, s), 8.08 (2H, d, J=8.4 Hz), 8.66 (1H, s), 10.41 (1H, s).


FABMS(pos) 418.2 [M+H]⁺

Melting point: 220 - 222°C.

10

Example 135

4'-Chloro-N-[6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



15

The titled compound was obtained by carrying out the same operation as in Example 51, using 4'-chloro-N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 56. ¹H-NMR (CDCl₃) δ: 2.30 (3H, s), 2.25-2.50 (10H, m), 2.83 (2H, t, J = 8.1 Hz), 3.07 (2H, s), 6.35 (1H, s), 7.01 (1H, d, J = 8.1 Hz), 7.36 (1H, d, J = 7.8 Hz), 7.44 (2H, d, J = 8.4 Hz), 7.51 (1H, s), 7.55 (2H, d, J = 8.4 Hz), 7.66 (2H, d, J = 8.4 Hz), 7.84 (1H, s), 7.93 (2H, d, J = 8.4 Hz). Melting point: 220 - 222°C (crystallization solvent:

25

tetrahydrofuran - n-hexane)

Example 136


4'-Chloro-N-[6-[[methyl[2-(1-piperidinyl)ethyl]amino]methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

30

WO 01/21577

PCT/JP00/06375

247



The titled compound was obtained by carrying out the same operation as in Example 51, using 4'-chloro-N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-


biphenyl]-4-carboxamide obtained in Reference Example 56.

¹H-NMR (CDCl₃) δ : 1.72-1.77 (6H, m), 2.25-2.36 (2H, m), 2.27 (3H, s), 2.52-2.63 (8H, m), 2.84 (2H, t, J = 8.0 Hz), 3.08 (2H, s), 6.35 (1H, s), 7.01 (1H, d, J = 8.1 Hz), 7.38 (1H, d, J = 8.1 Hz), 7.44 (2H, d, J = 8.4 Hz), 7.49 (1H, s), 7.55 (2H, d, J = 8.4 Hz), 7.66 (2H, d, J = 8.4 Hz), 7.83 (1H, s), 7.93 (2H, d, J = 8.4 Hz).

Melting point: 165 - 167°C (crystallization solvent : tetrahydrofuran - n-hexane)

Example 137

4'-Chloro-N-[6-[[methoxy(methyl)amino]methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 51, using 4'-chloro-N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-

biphenyl]-4-carboxamide obtained in Reference Example 56.

¹H-NMR (CDCl₃) δ : 2.41 (2H, t, J = 8.1 Hz), 2.61 (3H, s), 2.86 (2H, t, J = 8.1 Hz), 3.37 (2H, s), 3.52 (3H, s), 6.39 (1H, s), 7.03 (1H, d, J = 8.1 Hz), 7.36 (1H, d, J = 8.1 Hz), 7.44 (2H, d, J = 8.4 Hz), 7.53 (1H, s), 7.55 (2H, d, J = 8.4 Hz), 7.66 (2H, d, J = 8.4 Hz), 7.83 (1H, s), 7.93 (2H, d, J = 8.4 Hz).

WO 01/21577


PCT/JP00/06375

248

Melting point: 190 - 192°C (crystallization solvent :
ethyl acetate - n-hexane)

Example 138

5 4'-Chloro-N-[6-[[4-(1-piperidinyl)-1-piperidinyl]methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide




The titled compound was obtained by carrying out the
10 same operation as in Example 51, using 4'-chloro-N-[6-(chloromethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 56.
¹H-NMR (CDCl₃) δ : 1.45-1.96 (12H, m), 2.29-2.34 (3H, m),
2.57 (4H, s), 2.83 (2H, t, J = 8.1 Hz), 2.96-3.03 (4H, m),
15 6.32 (1H, s), 7.00 (1H, d, J = 8.1 Hz), 7.38 (1H, d, J = 8.1 Hz), 7.44 (2H, d, J = 8.4 Hz), 7.50 (1H, s), 7.55 (2H, d, J = 8.4 Hz), 7.66 (2H, d, J = 8.4 Hz), 7.86 (1H, s), 7.93 (2H, d, J = 8.4 Hz).

Melting point: 232 - 234°C (crystallization solvent :
20 ethyl acetate - n-hexane)

Example 139

6-(4-Fluorophenyl)-N-[3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl]nicotineamide



25 The titled compound was obtained by carrying out the same operation as in Example 1, using 3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine obtained in Reference Example 87.

WO 01/21577

PCT/JP00/06375


249

¹H-NMR (CDCl₃) δ : 1.70(4H, s), 2.43 (4H, s), 3.12 (2H, s), 4.73 (2H, s), 6.37 (1H, s), 7.03 (1H, d, J = 7.8 Hz), 7.29-7.40 (4H, m), 8.15 (1H, d, J = 8.4 Hz), 8.22-8.39 (3H, m), 9.15 (1H, s), 10.40 (1H, s).

5 Melting point: 233 - 235°C (crystallization solvent : tetrahydrofuran - n-hexane)

Example 140

4-Bromo-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]benzamide




The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ : 1.79 (4H, s), 2.35 (2H, t, J = 8.1 Hz), 2.52 (4H, s), 2.83 (2H, t, J = 8.1 Hz), 3.17 (2H, s), 6.35 (1H, s), 6.99 (1H, d, J = 8.1 Hz), 7.34 (1H, d, J = 8.1 Hz), 7.43 (1H, s), 7.60 (2H, d, J = 8.4 Hz), 7.72 (2H, d, J = 8.4 Hz), 7.76 (1H, s).

Melting point: 135 - 137°C (crystallization solvent : ethyl acetate - n-hexane)

Example 141

6-(4-Methoxyphenyl)-N-[3-(1-pyrrolidinymethyl)-2H-chromen-7-yl]nicotinamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 3-(1-pyrrolidinymethyl)-2H-chromen-7-ylamine

WO 01/21577

PCT/JP00/06375

250

pyrrolidinylmethyl)-2H-chromen-7-amine obtained in Reference Example 87.

¹H-NMR (CDCl₃) δ : 1.70 (4H, s), 2.44 (4H, s), 3.12 (2H, s), 3.84 (3H, s), 4.73 (2H, s), 6.37 (1H, s), 7.03 (1H, d, J = 8.1 Hz), 7.09 (2H, t, J = 8.7 Hz), 7.29 (1H, d, J = 8.4 Hz), 7.31 (1H, s), 8.07 (1H, d, J = 8.7 Hz), 8.16 (2H, d, J = 8.7 Hz), 8.32 (1H, d, J = 8.4 Hz), 9.12 (1H, s), 10.34 (1H, s).

Elemental analysis for C₂₇H₂₇N₃O₃


Calcd.: C, 73.45; H, 6.16; N, 9.52.

Found: C, 73.02; H, 6.27; N, 9.33.

Melting point: 243 - 245°C (crystallization solvent : tetrahydrofuran - n-hexane)

Example 142

4-(4-Fluorophenyl)-N-[3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine obtained in Reference Example 87.

¹H-NMR (CDCl₃) δ : 1.69- 1.91 (8H, m), 2.49 (4H, s), 2.70 (1H, t, J = 12.0 Hz), 2.97 (2H, t, J = 12.0 Hz), 3.12 (2H, s), 4.19 (2H, d, J = 13.0 Hz), 4.76 (2H, s), 6.26 (1H, s), 6.37 (1H, s), 6.82-7.03 (5H, m), 7.16 (2H, dd, J = 5.4, 8.4 Hz).

Melting point: 176 - 178°C (crystallization solvent : ethyl acetate - diisopropyl ether)

30

Example 143


N-[3-(1-Pyrrolidinylmethyl)-2H-chromen-7-yl][1,1'-

WO 01/21577

PCT/JP00/06375

251

biphenyl]-4-carboxamide




The titled compound was obtained by carrying out the same operation as in Example 1, using 3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine obtained in Reference Example 87.

¹H-NMR (CDCl₃) δ : 1.79 (4H, s), 2.50 (4H, s), 3.15 (2H, s), 4.81 (2H, s), 6.30 (1H, s), 6.95 (1H, d, J = 8.1 Hz), 7.13 (1H, s), 7.20 (1H, d, J = 8.1 Hz), 7.39-7.50 (3H, m), 7.61-7.70 (4H, m), 7.82 (1H, s), 7.92 (2H, d, J = 8.1 Hz).
Melting point: 198 - 200°C (crystallization solvent : ethyl acetate)

Example 144

N-[6-[(N-Benzyl-N-methylamino)methyl]-7,8-dihydro-2-naphthalenyl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-[(N-benzyl-N-methylamino)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 88.

¹H-NMR (CDCl₃) δ : 2.20 (3H, s), 2.38 (2H, t, J = 8.1 Hz), 2.85 (2H, t, J = 8.1 Hz), 3.09 (2H, s), 3.52 (2H, s), 6.39 (1H, s), 7.02 (1H, d, J = 8.1 Hz), 7.13-7.66 (13H, m), 7.84 (1H, s), 7.93 (2H, d, J = 8.4 Hz).

Melting point: 143 - 145°C (crystallization solvent : ethyl acetate - n-hexane)

Example 145


4'-Isobutyrylamino-N-[6-(1-pyrrolidinylmethyl)-7,8-

WO 01/21577

PCT/JP00/06375

252


dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as an amorphous powder by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54. MS m/z 494.4 (MH⁺).

Example 146


Ethyl 4'-[[[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]amino]carbonyl][1,1'-biphenyl]-3-carboxylate



The titled compound was obtained as an amorphous powder by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8--dihydro-2-naphthalenamine obtained in Reference Example 54. MS m/z 481.4 (MH⁺).

Example 147

3-[4'-[[[6-(1-Pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]amino]carbonyl][1,1'-biphenyl]-4-yl]propionic acid



The titled compound was obtained as a powder by carrying out the same operation as in Example 1, using

WO 01/21577

PCT/JP00/06375


253

6-(1-pyrrolidinymethyl)-7,8--dihydro-2-naphthalenamine
obtained in Reference Example 54.

MS m/z 481.4 (MH⁺).

5 Example 148

4'-Methoxy-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



10 The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.80 (4H, m), 2.36 (2H, t, J=7.8 Hz), 2.52 (4H, m), 2.86 (2H, t, J=7.8 Hz), 3.18 (2H, s), 3.87 (3H, s), 6.36 (1H, s), 7.00-7.03 (3H, m), 7.26 (1H, m), 7.38 (1H, d, J=8.3 Hz), 7.49 (1H, s), 7.58 (2H, d, J=8.6 Hz), 7.67 (1H, d, J=8.2 Hz), 7.78 (1H, s), 7.90 (2H, d, J=8.2 Hz).

Elemental analysis for C₂₉H₃₀N₂O₂


Calcd.: C, 79.42; H, 6.89; N, 6.39.

20 Found: C, 79.21; H, 6.88; N, 6.35.

Melting point: 187-188 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 149

25 6-(4-Fluorophenyl)-N-[6-[(1-pyrrolidinyl)methyl]-7,8-dihydro-2-naphthalenyl]nicotinamide



30 The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine

WO 01/21577

PCT/JP00/06375

254

obtained in Reference Example 54.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.81 (4H, m), 2.36 (2H, t, $J=8.1$ Hz),
2.53 (4H, m), 2.86 (2H, t, $J=8.1$ Hz), 3.18 (2H, s), 6.37
(1H, s), 7.03 (1H, d, $J=7.8$ Hz), 7.16-7.30 (3H, m), 7.47
(1H, s), 7.77-7.82 (2H, m), 8.06 (2H, dd, $J=8.9, 5.3$ Hz),
8.25 (1H, dd, $J=8.4, 2.2$ Hz), 9.11 (1H, d, $J=2.0$ Hz).

Elemental analysis for $\text{C}_{27}\text{H}_{26}\text{FN}_3\text{O}$


Calcd.: C, 75.85; H, 6.13; N, 9.83.

Found: C, 75.71; H, 5.93; N, 9.75.

Melting point: 225-227 °C (crystallization solvent:
ethyl acetate - diisopropyl ether)

Example 150

6-(4-Methylphenyl)-N-[6-[(1-pyrrolidinyl)methyl]-7,8-
dihydro-2-naphthalenyl]nicotinamide



The titled compound was obtained by carrying out the
same operation as in Example 1, using 6-(1-
pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
obtained in Reference Example 54.

$^1\text{H-NMR}$ (CDCl_3) δ : 1.81 (4H, m), 2.36 (2H, t, $J=7.8$ Hz), 2.43
(3H, s), 2.53 (4H, m), 2.86 (2H, t, $J=7.8$ Hz), 3.19 (2H,
s), 6.37 (1H, s), 7.02 (1H, d, $J=8.7$ Hz), 7.25-7.39 (3H,
m), 7.47 (1H, s), 7.82 (2H, m), 7.96 (2H, d, $J=8.1$ Hz), 8.23
(1H, dd, $J=8.1, 2.3$ Hz), 9.12 (1H, d, $J=2.3$ Hz).

Melting point: 235-236 °C (crystallization solvent: ethyl
acetate - diisopropyl ether)


Example 151

N-[6-[(Dimethylamino)methyl]-7,8-dihydro-2-
naphthalenyl]-6-(4-fluorophenoxy)nicotinamide

WO 01/21577

PCT/JP00/06375

255



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine
5 obtained in Reference Example 41-2).

¹H-NMR (CDCl₃) δ: 2.25 (6H, s), 2.34 (2H, t, J=8.1 Hz), 2.86 (2H, t, J=8.1 Hz), 2.99 (2H, s), 6.35 (1H, s), 7.03 (1H, d, J=8.1 Hz), 7.17 (2H, m), 7.26 (1H, m), 7.39 (1H, d, J=8.1 Hz), 7.47 (1H, s), 7.78 (1H, d, J=7.2 Hz), 7.83
10 (1H, s), 8.06 (1H, dd, J=8.4, 6.7 Hz), 8.25 (1H, d, J=6.7 Hz), 9.12 (1H, s).

Elemental analysis for C₂₅H₂₄FN₃O


Calcd.: C, 74.79; H, 6.03; N, 10.47.

Found: C, 74.74; H, 5.95; N, 10.24.

15 Melting point: 216-219 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 152

6-(2,4-Difluorophenyl)-N-[6-[(dimethylamino)methyl]-
20 7,8-dihydro-2-naphthalenyl]nicotinamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine
25 obtained in Reference Example 41-2).

¹H-NMR (CDCl₃) δ: 2.25 (6H, s), 2.34 (2H, t, J=8.1 Hz), 2.85 (2H, t, J=8.1 Hz), 3.00 (2H, s), 6.35 (1H, s), 6.90-7.06 (3H, m), 7.39 (1H, d, J=7.8 Hz), 7.47 (1H, s), 7.80-7.90 (2H, m), 8.10 (1H, dd, J=15.3, 8.8 Hz), 8.23 (1H, dd, J=8.4,
30 2.3 Hz), 9.15 (1H, d, J=1.7 Hz).

WO 01/21577

PCT/JP00/06375

256

Elemental analysis for $C_{25}H_{23}F_2N_3O$


Calcd.: C, 71.58; H, 5.53; N, 10.02.

Found: C, 71.50; H, 5.49; N, 9.61.

Melting point: 162-163 °C (crystallization solvent: ethyl
acetate - diisopropyl ether)

Example 153

6-Phenyl-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide



10

The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

1H -NMR ($CDCl_3$) δ : 1.81 (4H, m), 2.36 (2H, t, J=8.1 Hz), 2.53 (4H, m), 2.85 (2H, t, J=8.1 Hz), 3.18 (2H, s), 6.37 (1H, s), 7.02 (1H, d, J=8.1 Hz), 7.37-7.53 (5H, m), 7.83 (1H, d, J=8.1 Hz), 7.86 (1H, d, J=6.2 Hz), 8.04 (1H, s), 8.06 (1H, d, J=1.7 Hz), 8.24 (1H, dd, J=8.4, 2.4 Hz), 9.13 (1H, d, J=2.2 Hz).

20

Elemental analysis for $C_{27}H_{27}N_3O$

Calcd.: C, 79.19; H, 6.65; N, 10.26.


Found: C, 78.93; H, 6.65; N, 10.19.

Melting point: 186-187 °C (crystallization solvent: ethyl
acetate - diisopropyl ether)

25

Example 154

6-(4-Methoxyphenyl)-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide



30

WO 01/21577

PCT/JP00/06375


257

The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

- 5 ¹H-NMR (CDCl₃) δ: 1.80 (4H, m), 2.36 (2H, t, J=8.1 Hz), 2.52 (4H, m), 2.84 (2H, t, J=8.1 Hz), 3.18 (2H, s), 3.88 (3H, s), 6.36 (1H, s), 7.02 (3H, m), 7.37 (1H, d, J=7.5 Hz), 7.47 (1H, s), 7.78 (1H, d, J=8.1 Hz), 7.79 (1H, s), 8.03 (2H, d, J=8.5 Hz), 8.20 (1H, d, J=8.1 Hz), 9.08 (1H, s).
- 10 Melting point: 219-220 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 155

- 15 4-(4-Methylphenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

- 20 ¹H-NMR (CDCl₃) δ: 1.64-1.92 (8H, m), 2.29 (2H, m), 2.32 (3H, s), 2.51 (4H, m), 2.64 (1H, m), 2.80 (2H, t, J=7.8 Hz), 2.97 (2H, dd, J=13.1, 10.7 Hz), 3.15 (2H, s), 4.19 (2H, d, J=13.1 Hz), 6.32 (1H, s), 6.35 (1H, s), 6.42 (1H, d, J=7.8 Hz), 7.06-7.20 (6H, m)

Elemental analysis for C₂₈H₃₅N₃O · 0.5H₂O

Calcd.: C, 76.67; H, 8.27; N, 9.58.

Found: C, 76.72; H, 8.03; N, 9.36.

- 25 Melting point: 197-198 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 156


4-Phenyl-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-

WO 01/21577

PCT/JP00/06375

258

naphthalenyl]-1-piperidinecarboxamide




The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.72-1.94 (8H, m), 2.32 (2H, t, J=8.1 Hz), 2.50 (4H, m), 2.72 (1H, m), 2.80 (2H, t, J=8.1 Hz), 2.99 (2H, dd, J=13.4, 10.6 Hz), 3.16 (2H, s), 4.21 (2H, d, J=13.4 Hz), 6.32 (1H, s), 6.34 (1H, s), 6.93 (1H, d, J=8.4 Hz), 7.07 (1H, d, J=8.1 Hz), 7.20-7.35 (6H, m).

Melting point: 184-186 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 157

4-(1,3-Benzodioxol-5-yl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.61-1.88 (8H, m), 2.31 (2H, t, J=8.1 Hz), 2.51 (4H, m), 2.59 (1H, m), 2.62 (2H, t, J=8.1 Hz), 2.94 (2H, dd, J=13.1, 11.2 Hz), 3.15 (2H, s), 4.18 (2H, d, J=13.1 Hz), 5.93 (2H, s), 6.31 (1H, s), 6.44 (1H, s), 6.64-6.77 (3H, m), 6.92 (1H, d, J=8.1 Hz), 7.07 (1H, d, J=8.1 Hz), 7.19 (1H, s).

Melting point: 149-150 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


WO 01/21577

PCT/JP00/06375

259

Example 158

4-(4-Fluorophenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-
dihydro-2-naphthalenyl]-3,6-dihydro-1(2H)-
5 pyridinecarboxamide



The titled compound was obtained by carrying out the
same operation as in Example 99, using 6-(1-
pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
10 obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.79 (4H, m), 2.32 (2H, t, J=8.1 Hz), 2.50
(4H, m), 2.59 (2H, brt), 2.80 (2H, t, J=8.1 Hz), 3.17 (2H,
s), 3.74 (2H, t, J=5.7 Hz), 4.15 (2H, d, J=2.5 Hz), 6.00
(1H, brt), 6.32 (1H, s), 6.32 (1H, s), 6.94 (1H, d, J=8.1
15 Hz), 7.00-7.32 (6H, m).

Elemental analysis for C₂₇H₃₀FN₃O


Calcd.: C, 75.15; H, 7.01; N, 9.74.

Found: C, 75.09; H, 6.93; N, 9.77.

Melting point: 206-207 °C (crystallization solvent: ethyl
20 acetate - diisopropyl ether)

Example 159

4-(4-Chlorophenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-
dihydro-2-naphthalenyl]-3,6-dihydro-1(2H)-
25 pyridinecarboxamide



The titled compound was obtained by carrying out the
same operation as in Example 99, using 6-(1-
pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
30 obtained in Reference Example 54.

WO 01/21577

PCT/JP00/06375

260

¹H-NMR (CDCl₃) δ : 1.79 (4H, m), 2.32 (2H, t, J=8.1 Hz), 2.50 (4H, m), 2.59 (2H, brt), 2.80 (2H, t, J=8.1 Hz), 3.16 (2H, s), 3.73 (2H, t, J=5.6 Hz), 4.15 (2H, d, J=2.8 Hz), 6.06 (1H, brt), 6.30 (1H, s), 6.32 (1H, s), 6.93 (1H, d, J=7.8 Hz), 7.09 (1H, d, J=7.8 Hz), 7.21-7.31 (5H, m).

Elemental analysis for C₂₇H₃₀ClN₃O


Calcd.: C, 72.39; H, 6.75; N, 9.38.

Found: C, 72.19; H, 6.75; N, 9.19.

Melting point: 217-218 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 160

4-(4-Chlorophenyl)-4-hydroxy-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ : 1.79 (4H, m), 1.80 (2H, m), 2.04 (1H, dd, J=13.1, 10.8 Hz), 2.06 (1H, dd, J=13.1, 10.8 Hz), 2.31 (2H, t, J=7.8 Hz), 2.50 (1H, brs), 2.51 (4H, m), 2.79 (2H, t, J=7.8 Hz), 3.15 (2H, s), 3.41 (2H, dd, J=12.6, 10.8 Hz), 4.00 (2H, d, J=12.6 Hz), 6.32 (1H, s), 6.37 (1H, s), 6.93 (1H, d, J=8.1 Hz), 7.05-7.42 (6H, m).

Melting point: 181-182 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


Example 161

4-(4-Methylphenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-3,6-dihydro-1(2H)-pyridinecarboxamide

WO 01/21577

PCT/JP00/06375

261



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine


5 obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ: 1.79 (4H, m), 2.32 (2H, t, J=7.8 Hz), 2.35 (3H, s), 2.50 (4H, m), 2.61 (2H, brt), 2.80 (2H, t, J=7.8 Hz), 3.16 (2H, s), 3.73 (2H, t, J=5.7 Hz), 4.15 (2H, d, J=2.8 Hz), 6.03 (1H, s), 6.29 (1H, s), 6.32 (1H, s), 6.93 (1H, d, J=8.1 Hz), 7.07-7.30 (6H, m).

Melting point: 199-202 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 162

15 6-(4-Chlorophenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine

20 obtained in Reference Example 54.

¹H-NMR (CDCl₃+DMSO-d₆) δ: 1.80 (4H, m), 2.32-2.58 (6H, m), 2.85 (2H, t, J=8.0 Hz), 3.18 (2H, s), 6.36 (1H, s), 7.01 (1H, d, J=8.4 Hz), 7.48 (2H, d, J=8.4 Hz), 7.49 (1H, m), 7.59 (1H, s), 7.83 (1H, d, J=8.4 Hz), 8.04 (2H, d, J=8.4 Hz), 8.35 (1H, dd, J=8.4, 2.2 Hz), 9.25 (1H, d, J=2.2 Hz), 9.42 (1H, s).

Elemental analysis for C₂₇H₂₆ClN₃O

Calcd.: C, 73.04; H, 5.90; N, 9.46.

30 Found: C, 73.11; H, 5.71; N, 9.20.

WO 01/21577


PCT/JP00/06375

262

Melting point: 252-253 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 163

5 N-[6-[(Dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl]-6-(4-methylphenyl)nicotinamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 6-[(N,N-
10 dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 41-2).


¹H-NMR (CDCl₃) δ: 2.25 (6H, s), 2.34 (2H, t, J=8.1 Hz), 2.43 (3H, s), 2.85 (2H, t, J=8.1 Hz), 2.99 (2H, s), 6.34 (1H, s), 7.02 (1H, d, J=8.1 Hz), 7.31 (2H, d, J=8.1 Hz),
15 7.39 (1H, d, J=8.1 Hz), 7.46 (1H, s), 7.81 (1H, d, J=8.4 Hz), 7.87 (1H, s), 7.96 (2H, d, J=8.1 Hz), 8.22 (1H, dd, J=8.4, 2.3 Hz), 9.11 (1H, d, J=2.3 Hz).

Melting point: 228-230 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

20

Example 164

6-(4-Chlorophenyl)-N-[6-[(dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl]nicotinamide



25 The titled compound was obtained by carrying out the same operation as in Example 1, using 6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 41-2).

¹H-NMR (CDCl₃) δ: 2.25 (6H, s), 2.35 (2H, t, J=8.1 Hz), 2.86 (2H, t, J=8.1 Hz), 2.99 (2H, s), 6.35 (1H, s), 7.04 (1H,
30

WO 01/21577

PCT/JP00/06375

263

d, J=8.1 Hz), 7.40 (1H, d, J=8.4 Hz), 7.49 (1H, brs), 7.49 (2H, d, J=8.4 Hz), 7.78 (1H, s), 7.84 (1H, d, J=8.4 Hz), 8.02 (2H, d, J=8.4 Hz), 8.26 (1H, dd, J=8.1, 2.2 Hz), 9.13 (1H, d, J=2.2 Hz).

5 Elemental analysis for $C_{25}H_{24}ClN_3O$

Calcd.: C, 71.85; H, 5.79; N, 10.05.


Found: C, 71.88; H, 5.67; N, 9.86.

Melting point: 248-249 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

10

Example 165

4-(4-Chlorophenyl)-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



15

The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

1H -NMR ($CDCl_3$) δ : 1.66-1.91 (8H, m), 2.32 (2H, t, J=8.1 Hz), 2.50 (4H, m), 2.70 (1H, m), 2.80 (2H, t, J=8.1 Hz), 2.98 (2H, dd, J=13.7, 12.0 Hz), 3.16 (2H, s), 4.20 (2H, d, J=13.7 Hz), 6.32 (1H, s), 6.32 (1H, s), 6.93 (1H, d, J=8.1 Hz), 7.05-7.30 (6H, m).

20

Elemental analysis for $C_{27}H_{32}ClN_3O$

25

Calcd.: C, 72.06; H, 7.17; N, 9.34.

Found: C, 72.08; H, 7.23; N, 9.15.

Melting point: 194-195 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

30


Example 166

N-[6-[(Dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl]-4-(4-fluorophenyl)-1-piperidinecarboxamide

WO 01/21577

PCT/JP00/06375

264



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine


5 obtained in Reference Example 41-2).

¹H-NMR (CDCl₃) δ: 1.65-1.75 (2H, m), 1.89 (2H, d, J=11.4 Hz), 2.23 (6H, s), 2.30 (2H, t, J=8.1 Hz), 2.70 (1H, m), 2.80 (2H, t, J=8.1 Hz), 2.94-3.01 (4H, m), 4.20 (2H, d, J=13.4 Hz), 6.30 (1H, s), 6.35 (1H, s), 6.92-7.20 (7H, m).

10 Melting point: 187-188 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 167

N-[6-[(Dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl]-4-(4-methylphenyl)-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine

20 obtained in Reference Example 41-2).

¹H-NMR (CDCl₃) δ: 1.66-1.74 (2H, m), 1.89 (2H, d, J=11.7 Hz), 2.28 (6H, s), 2.30 (2H, t, J=8.1 Hz), 2.38 (3H, s), 2.68 (1H, m), 2.80 (2H, t, J=8.1 Hz), 2.94-3.02 (4H, m), 4.19 (2H, d, J=12.8 Hz), 6.30 (1H, s), 6.35 (1H, s), 6.93 (1H, d, J=8.1 Hz), 7.07-7.20 (6H, m).

Elemental analysis for C₂₆H₃₃N₃O · 0.5H₂O

Calcd.: C, 75.69; H, 8.31; N, 10.18

Found: C, 75.44; H, 8.16; N, 10.05

30 Melting point: 200-202 °C (crystallization solvent: ethyl

WO 01/21577


PCT/JP00/06375

265

acetate - diisopropyl ether)

Example 168

N-[6-[(Dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-2-carboxamide
hydrochloride




The titled compound was obtained by carrying out the same operation as in Example 1, using 6-amino-2-[(N,N-dimethylamino)methyl]tetralin hydrochloride.

¹H-NMR (DMSO-d₆) δ: 1.39 (1H, m), 1.99 (1H, m), 2.17 (1H, m), 2.42 (1H, dd, J=16.2, 10.1 Hz), 2.78 (6H, s), 2.88 (1H, dd, J=16.2, 4.5 Hz), 3.06 (2H, t, J=5.7 Hz), 3.38 (2H, s), 6.94-7.62 (11H, m), 7.64 (1H, d, J=1.7 Hz), 10.11 (1H, brs), 10.18 (1H, s).

Melting point: 196-197 °C (crystallization solvent: methanol - ethyl acetate)

Example 169

N-[6-[(Dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide hydrochloride



4'-Fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide synthesized in Example 42 was dissolved in ethyl acetate. An excess amount of 4N hydrochloric acid-ethyl acetate solution was added to the solution, which was concentrated under reduced pressure. The resulting residue was recrystallized from methanol - ethyl

WO 01/21577

PCT/JP00/06375

266

acetate, to give the titled compound.


¹H-NMR (DMSO-d₆) δ: 1.43 (1H, m), 2.06 (1H, m), 2.21 (1H, m), 2.45 (1H, m), 2.79 (6H, s), 2.92 (1H, dd, J=16.2, 4.2 Hz), 3.08 (2H, d, J=6.4 Hz), 3.33 (2H, s), 7.05 (1H, d, J=8.4 Hz), 7.34 (2H, dd, J=8.9, 8.9 Hz), 7.53 (1H, d, J=8.4 Hz), 7.59 (1H, s), 7.80 (4H, m), 8.06 (2H, d, J=8.1 Hz), 10.02 (1H, s), 10.03 (1H, brs).

Melting point: 240-245 °C (crystallization solvent: methanol - ethyl acetate)

10

Example 170

6-(4-Fluorophenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide hydrochloride



15

The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.


¹H-NMR (DMSO-d₆) δ: 1.70 (4H, m), 2.26 (2H, t, J=8.1 Hz), 2.44 (4H, m), 2.76 (2H, t, J=8.1 Hz), 3.12 (2H, s), 3.34 (1H, s), 6.36 (1H, s), 7.03 (1H, d, J=7.8 Hz), 7.37 (2H, dd, J=8.4, 7.0 Hz), 7.57 (1H, d, J=8.4 Hz), 7.59 (1H, s), 8.13-8.42 (4H, m), 9.19 (1H, s), 10.43 (1H, s).

Melting point: 229-231 °C (crystallization solvent: methanol - ethyl acetate)

25

Example 171

6-(4-Fluorophenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide dihydrochloride



30

WO 01/21577

PCT/JP00/06375


267

The titled compound was obtained by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

- 5 $^1\text{H-NMR}$ (DMSO-d_6) δ : 2.00 (4H, m), 2.45 (4H, m), 2.83 (2H, t, $J=8.1$ Hz), 3.05 (2H, m), 3.47 (2H, m), 3.88 (1H, s), 6.69 (1H, s), 7.13 (1H, d, $J=8.1$ Hz), 7.38 (2H, dd, $J=8.9$, 8.6 Hz), 7.64 (1H, d, $J=10.6$ Hz), 7.66 (1H, s), 8.14-8.42 (4H, m), 9.19 (1H, s), 10.52 (1H, s), 10.60 (1H, brs).
- 10 Melting point: 245-248 $^{\circ}\text{C}$ (crystallization solvent: methanol - ethyl acetate)

Example 172

- 15 N-[6-[(Dimethylnitroyl)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide 3-chlorobenzoate



- 4'-FluoroN-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide (100 mg) obtained in Example 42 was dissolved in acetone (10 ml), which was stirred under ice-cooling.
- 20 3-Chloroperbenzoic acid (purity : 50%) (86 mg) was added to the solution, which was stirred under ice-cooling for 1 hour. The reaction mixture was concentrated under reduced pressure, and the residue was washed with diisopropyl ether, to give the titled compound (158 mg).
- 25 $^1\text{H-NMR}$ (DMSO-d_6) δ : 1.57 (1H, m), 2.07 (1H, m), 2.61 (1H, m), 2.82 (2H, m), 3.04 (1H, m), 3.33 (1H, m), 3.48 (6H, s), 3.56-3.67 (2H, m), 6.55 (1H, s), 7.03 (1H, d, $J=8.4$ Hz), 7.30-7.56 (6H, m), 7.78-7.85 (6H, m), 8.04 (2H, d, $J=8.4$ Hz), 10.17 (1H, s).
- 30

WO 01/21577


PCT/JP00/06375

268

FABMS(pos) 419.1 [M+H]⁺

Example 173

N-[6-[(Dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-sulfonamide
hydrochloride



6-[(N, N-Dimethylamino)methyl]-7,8-dihydro-2-naphthalenamine (200 mg, 0.72 mmol) obtained in Example 41-2) was dissolved in acetonitrile (30 ml). Triethylamine (0.401 ml, 2.88 mmol) and [1,1'-biphenyl]-4-sulfonylchloride (200 mg, 0.79 mmol) were added to the solution under ice-cooling, which was stirred for 3 hours. The reaction mixture was concentrated. Ethyl acetate and water were added to the residue, and extraction was conducted. The ethyl acetate layer was concentrated, and the residue was purified by alumina column chromatography (development solvent; ethyl acetate:n-hexane = 33:67). 4N Hydrogen chloride-ethyl acetate solution was added to the resulting oily substance, which was concentrated. The residue was recrystallized from methanol - ethyl acetate, to give the titled compound (194 mg).


¹H-NMR (DMSO-d₆) δ: 1.32 (1H, m), 1.96 (1H, m), 2.11 (1H, m), 2.35 (1H, d, J=15.9, 10.0 Hz), 2.74 (2H, m), 2.78 (7H, m), 3.02 (2H, m), 6.89 (2H, d, J=10.6 Hz), 6.91 (1H, m), 7.40-7.51 (3H, m), 7.70 (2H, d, J=6.7 Hz), 7.85 (4H, m), 9.92 (1H, brs), 10.23 (1H, s).

Melting point: 168-170 °C (crystallization solvent: methanol - ethyl acetate)

FABMS(pos) 421.1 [M+H]⁺

Example 174

4'-Chloro-N-[4-(4-piperidinyl)phenyl][1,1'-biphenyl]-4-carboxamide




5 The titled compound was obtained as a colorless powder by carrying out the same operation as in Example 127-2), using 4'-chloro-N-[4-(4-piperidinyl)phenyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 89. ¹H-NMR (CDCl₃+ DMSO-d₆) δ: 1.40-1.90 (4H, m), 2.60-2.90 (3H, m), 3.18-3.28 (2H, m), 7.19 (2H, d, J=8.1 Hz), 7.49 (2H, d, J=7.0 Hz), 7.67-7.75 (6H, m), 8.07-8.10 (3H, m), 10.16 (1H, s).
Melting point: 276-281 °C (decomposition) (crystallization solvent: ethyl acetate)

15

Example 175

4'-Chloro-N-[4-(1-methyl4-piperidinyl)phenyl][1,1'-biphenyl]-4-carboxamide



20 A mixture of 4'-chloro-N-[4-(4-piperidinyl)phenyl][1,1'-biphenyl]-4-carboxamide (0.17 g) obtained in Example 174, 37% aqueous formaldehyde solution (0.05 ml) and formic acid (0.5 ml) was heated at 100°C for 4 hours. The reaction mixture was cooled to room
25 temperature. Water was added to the mixture, which was made alkaline with 8N aqueous sodium hydroxide solution, and extracted with ethyl acetate - tetrahydrofuran (1:1) mixed solution. The extract was washed with saturated aqueous sodium chloride solution, dried over anhydrous magnesium sulfate, and then the solvent was distilled out under
30 reduced pressure. The resulting solid was washed with ethyl acetate, dried under reduced pressure, to give the titled compound (90 mg).

¹H-NMR (CDCl₃+ DMSO-d₆) δ: 1.55-1.80 (2H, m), 1.90-2.10

WO 01/21577

PCT/JP00/06375

270


(2H, m), 2.22 (3H, s), 2.30-2.45 (1H, m), 2.80-3.20 (4H, m), 7.11 (2H, d, J=8.1 Hz), 7.36 (2H, d, J=8.1 Hz), 7.50-7.63 (6H, m), 7.97 (2H, d, J=8.4 Hz), 9.79 (1H, s).

Melting point: 273-277 °C (decomposition) (Washing

5 solvent: ethyl acetate)

Example 176

Benzyl 4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenylcarbamate



10

N,N-Dimethylethylenediamine (0.64 ml), WSC (1.31 g), HOBT (1.05 g), and triethylamine (2.4 ml) were added to a tetrahydrofuran (50 ml) solution of 2-[4-

15 [[(benzyloxy)carbonyl]amino]phenyl]acetic acid (1.5 g)


obtained in Reference Example 90. After stirring for 20 hours, the reaction mixture was poured into water, and extraction was conducted using ethyl acetate. The organic layer was washed with water, saturated aqueous sodium bicarbonate solution, and saturated aqueous sodium chloride solution, dried and then concentrated. The residue was recrystallized from ethyl acetate - hexane, to give the titled compound (1.72 g).

20

Melting point: 126-127 °C.

25 Example 177

N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride



30

Oxalyl chloride (0.56 ml) was added dropwise to a

tetrahydrofuran (45 ml) solution of 4-biphenylcarboxylic acid (1.01 g) under ice-cooling. 9 drops of DMF was added to the mixture, and the temperature of the mixture was raised to room temperature, which was stirred for 40 minutes. The reaction mixture was concentrated and dried.

A tetrahydrofuran (50 ml) solution of the residue was added dropwise to a tetrahydrofuran (45 ml) solution of 2-(4-aminophenyl)-N-[2-(dimethylamino)ethyl]acetamide (939 mg) obtained in Reference Example 91 under ice-cooling.

Then the temperature of the reaction mixture was raised to room temperature, which was stirred for 2 hours. Saturated aqueous sodium bicarbonate solution was added to the reaction mixture, and extraction was conducted using ethyl acetate. The organic layer was washed with water and saturated aqueous sodium chloride solution, dried over sodium sulfate, and then concentrated. The residue was dissolved in tetrahydrofuran. 4N Hydrochloric acid-ethyl acetate was added to the solution, which was concentrated.

The residue was recrystallized from methanol - diisopropyl ether, to give the titled compound (750 mg). Melting point: 216-217 °C.

The above N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl][1,1'-biphenyl]-4-carboxamide hydrochloride (100 mg) was dissolved in saturated aqueous sodium bicarbonate solution, and extraction was conducted using tetrahydrofuran-ethyl acetate (1:1). The organic layer was washed with saturated aqueous sodium chloride solution, dried over sodium sulfate, and then concentrated.

The residue was recrystallized from methanol - diisopropyl ether, to give a free base form (56 mg) of the titled compound.

Melting point: 228-229 °C.


Example 178

Benzyl 4-[[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]anilino]carbonyl]-1-piperidinecarboxylate

WO 01/21577

PCT/JP00/06375


272



2-(4-Aminophenyl)-N-[2-(dimethylamino)ethyl]acetamide (221 mg), WSC (249 mg), 1-hydroxybenzotriazole (199 mg), triethylamine (0.4 ml), and dimethylaminopyridine (244 mg) were added to a tetrahydrofuran (10 ml) solution of 1-[(benzyloxy)carbonyl]-4-piperidinecarboxylic acid (290 mg), which was stirred for 20 hours. The reaction mixture was poured into water, and extraction was conducted using ethyl acetate. The organic layer was washed with water, saturated aqueous sodium bicarbonate solution, and saturated aqueous sodium chloride solution, dried over sodium sulfate, and then concentrated. The residue was recrystallized from methanol - diisopropyl ether, to give the titled compound (230 mg). Melting point: 169-170 °C.

Example 179

N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-3-[3-(2-naphthyl)-1,2,4-oxadiazol-5-yl]propanamide




2-(4-Aminophenyl)-N-[2-(dimethylamino)ethyl]acetamide (221 mg), WSC (249 mg), 1-hydroxybenzotriazole (199 mg), triethylamine (0.4 ml), and dimethylaminopyridine (244 mg) were added to a DMF (5 ml) solution of 3-[3-(2-naphthyl)-1,2,4-oxadiazol-5-yl]propionic acid (268 mg), which was stirred for 5 hours. The reaction mixture was poured into water, and extraction was conducted using ethyl acetate. The organic layer was

washed with water, saturated aqueous sodium bicarbonate solution, and saturated aqueous sodium chloride solution, dried over sodium sulfate, and then concentrated. The residue was recrystallized from methanol, to give the
5 titled compound (166 mg).
Melting point: 173-174 °C.

Example 180

N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-2-(4-nitrophenyl)acetamide
10




2-(4-Aminophenyl)-N-[2-(dimethylamino)ethyl]acetamide (221 mg), WSC (free form : 0.23 ml), 1-hydroxybenzotriazole (199 mg), and
15 dimethylaminopyridine (244 mg) were added to a DMF (5 ml) solution of 4-nitrophenylacetic acid (181 mg), which was stirred for 4 hours. The reaction mixture was poured into water, and extraction was conducted using ethyl acetate.
The organic layer was washed with water, saturated aqueous
20 sodium bicarbonate solution, and saturated aqueous sodium chloride solution, dried over sodium sulfate, and then concentrated. The residue was recrystallized from methanol, to give the titled compound (80 mg).
Melting point: 160-162 °C.

25

Example 181

(E)-N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-3-[4-(4-methoxyphenoxy)phenyl]-2-propanamide




30

2-(4-Aminophenyl)-N-[2-(dimethylamino)ethyl]acetamide (221 mg), WSC (free form :

0.23 ml), 1-hydroxybenzotriazole (199 mg), triethylamine (0.14 ml) and dimethylaminopyridine (122 mg) were added to a DMF (5 ml) solution of (E)-3-[4-(4-methoxyphenoxy)phenyl]-2-propenoic acid (270 mg), which
5 was stirred for 24 hours. The reaction mixture was poured into water, and extraction was conducted using ethyl acetate - tetrahydrofuran (1:1). The organic layer was washed with water, saturated aqueous sodium bicarbonate solution, and saturated aqueous sodium chloride solution,
10 dried over sodium sulfate, and then concentrated. The resulting crude crystals were washed with diisopropyl ether, to give the titled compound (227 mg).
Melting point: 175-177 °C (decomposition).

15 Compounds described in the following Example 182 to 198 were produced in the same manner as in Example 181.
Example 182

4-[3-(1-Benzofuran-2-yl)-1,2,4-oxadiazol-5-yl]-N-[4-[2-
[[2-(dimethylamino)ethyl]amino]-2-
20 oxoethyl]phenyl]butanamide




Melting point: 161-163 °C.

Washing solvent: diisopropyl ether.

25 Example 183

N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-3-methoxy-4-(2-quinolinylmethoxy)benzamide



30 Melting point: 209-210 °C (decomposition).

Washing solvent: diisopropyl ether.


WO 01/21577

PCT/JP00/06375

275

Example 184

3-[1-(2,4-Dichlorobenzyl)-1H-indol-3-yl]-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]propanamide




5

Melting point: : 123-125 °C (decomposition).

Washing solvent: diisopropyl ether.

Example 185

10 N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-1-benzothiophen-2-carboxamide




Melting point: 186-187 °C (decomposition).

Washing solvent: diisopropyl ether.

15

Example 186

2-(2-Benzylphenyl)-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]acetamide



20 Melting point: 115-117 °C.

Washing solvent: diisopropyl ether.


Example 187

25 2-(3,4-dimethoxyphenyl)-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]acetamide

WO 01/21577

PCT/JP00/06375

276




Melting point: 123-124 °C.

Recrystallization solvent: methanol - diisopropyl ether.

5 Example 188

N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-2-(5-methoxy-2-methyl-1H-indol-3-yl)acetamide




10 Melting point: 125-126 °C.

Recrystallization solvent: methanol - diisopropyl ether.

Example 189

15 N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-4-(1H-indol-3-yl)butanamide




Melting point: 132-133 °C.

Washing solvent: diisopropyl ether.

20 Example 190

N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]furo[2,3-f][1,3]benzodioxol-6-carboxamide



WO 01/21577

PCT/JP00/06375


277

Melting point: : 173-175 °C (decomposition).

Washing solvent: diisopropyl ether.

Example 191

5 4-([1,1'-Biphenyl]-4-ylmethoxy)-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]benzamide




Melting point: 204-208 °C.

Washing solvent: diisopropyl ether.

10

Example 192

4-(Benzoylamino)-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]benzamide




15 Melting point: 220-221 °C.

Washing solvent: diisopropyl ether.

Example 193

20 N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-4'-methoxy[1,1'-biphenyl]-4-carboxamide



Melting point: 196-198 °C (decomposition).

Washing solvent: diisopropyl ether.

25


WO 01/21577

PCT/JP00/06375

278

Example 194

N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-9,10,10-trioxo-9,10-dihydro-10 λ ⁶-thioxanten-3-carboxamide




5

Melting point: : 162-163 °C (decomposition).

Washing solvent: diisopropyl ether.

Example 195

10 4-(Benzyloxy)-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]benzamide




Melting point: 190-192 °C (decomposition).

15 Washing solvent: diisopropyl ether.

Example 196

4-Benzoyl-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]benzamide



20

Melting point: 173-175 °C (decomposition).

Washing solvent: diisopropyl ether.


Example 197

25 N-[4-[2-[[2-(Dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-5-methyl-3-(4-pyridinyl)-1H-pyrrole-2-carboxamide

WO 01/21577

PCT/JP00/06375

279




Melting point: : 215-218 °C (decomposition).

Washing solvent : diisopropyl ether.

5 Example 198

1-(3,4-Dichlorobenzyl)-N-[4-[2-[[2-(dimethylamino)ethyl]amino]-2-oxoethyl]phenyl]-4-piperidinecarboxamide




10 Melting point: : 182-183 °C (decomposition).

Washing solvent : diisopropyl ether.

Example 199

15 4-(4-Methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

¹H-NMR (CDCl₃) δ : 1.61-1.91 (8H, m), 2.31 (2H, t, J=8.1 Hz), 2.54 (4H, m), 2.73-2.81 (3H, m), 2.98 (2H, t, J=7.8 Hz), 3.16 (2H, s), 3.79 (3H, s), 4.20 (2H, d, J=13.1 Hz), 6.31 (1H, s), 6.36 (1H, s), 6.86 (2H, d, J=8.6 Hz), 7.06-7.20 (5H, m).

Melting point: 175-176 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


WO 01/21577

PCT/JP00/06375

280

Example 200

4'-Methoxy-N-[6-(1-pyrrolidinylmethyl)-5-methyl-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 69.

10 ¹H-NMR (CDCl₃) δ: 1.78 (4H, m), 2.10 (3H, s), 2.37 (2H, t, J=8.1 Hz), 2.53 (4H, m), 2.76 (2H, t, J=8.1 Hz), 3.28 (2H, s), 3.87 (3H, s), 7.01 (2H, d, J=8.6 Hz), 7.27 (1H, d, J=7.8 Hz), 7.46 (1H, d, J=7.8 Hz), 7.48 (1H, s), 7.57 (2H, d, J=8.6 Hz), 7.66 (2H, d, J=8.4 Hz), 7.81 (1H, s), 7.92 (2H, d, J=8.4 Hz).

15 Elemental analysis for C₃₀H₃₂N₂O₂


Calcd.: C, 79.61; H, 7.13; N, 6.19

Found: C, 79.35; H, 7.28; N, 6.24

20 Melting point: 179-180 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 201

4-(4-Methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-5-methyl-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine

WO 01/21577

PCT/JP00/06375

281

obtained in Reference Example 69.

¹H-NMR (CDCl₃) δ: 1.67 (2H, dd, J=13.4, 4.0 Hz), 1.78 (4H, m), 1.89 (2H, d, J=11.4 Hz), 2.07 (3H, s), 2.34 (2H, t, J=7.5 Hz), 2.52 (4H, m), 2.68-2.73 (3H, m), 2.98 (2H, t, J=7.5 Hz),
5 3.26 (2H, s), 3.80 (3H, s), 4.20 (2H, d, J=13.4 Hz), 6.36 (1H, s), 6.86 (2H, d, J=8.4 Hz), 7.12-7.20 (5H, m).

Elemental analysis for C₂₈H₃₇N₃O₂


Calcd.: C, 75.13; H, 8.33; N, 9.39

Found: C, 74.96; H, 8.14; N, 9.10

10 Melting point: 163-164 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 202

4'-Fluoro-N-methyl-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide
15 hydrochloride



The titled compound was obtained by carrying out the same operation as in Example 1, using N-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
20 hydrochloride obtained in Reference Example 95.

¹H-NMR (DMSO-d₆) δ: 1.92-1.98 (4H, m), 2.39 (2H, t, J=8.1 Hz), 2.73 (2H, t, J=8.1 Hz), 3.00 (2H, m), 3.35 (3H, m), 3.44 (2H, m), 3.83 (2H, d, J=5.6 Hz), 6.62 (1H, s), 6.92-7.01
25 (2H, m), 7.11 (1H, s), 7.26 (2H, dd, J=8.9, 5.6 Hz), 7.38 (2H, d, J=8.1 Hz), 7.55 (2H, d, J=8.1 Hz), 7.69 (2H, dd, J=8.9, 5.6 Hz), 10.60 (1H, brs).

FABMS(pos) 441.2 [M+H]⁺


30 Example 203

N-[6-[(Dimethylamino)methyl]-5-hydroxy-5,6,7,8-tetrahydro-2-naphthalenyl]-4-(4-fluorophenyl)-1-piperidinecarboxamide

WO 01/21577

PCT/JP00/06375


282



N,N-Dimethylmethylene ammonium chloride (638 mg, 6.82 mmol) was added to a mixed solution of 4-(4-fluorophenyl)-N-(5-oxo-5,6,7,8-tetrahydro-2-naphthalenyl)-1-piperidinecarboxamide (1.00 g, 2.73 mmol) obtained in Reference Example 97 in tetrahydrofuran (10 ml) and acetonitrile (10 ml), which was stirred at room temperature for 1 day. The solvent was distilled out under reduced pressure. Ethyl acetate was added to the residue, which was washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. The resulting oily substance was dissolved in methanol (15 ml). Sodium borohydride (103 mg, 2.73 mmol) was added to the solution under ice-cooling, which was stirred for 1 hour. Then, the solvent was distilled out under reduced pressure. 1N Hydrochloric acid was added to the residue, which was washed with ethyl acetate. 4N Sodium hydroxide was added to the water layer to make it alkaline. The reaction mixture was extracted with ethyl acetate, which was washed with saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. The resulting residue was purified by aluminum B column chromatography (development solvent; ethyl acetate), powdered with hexane, to give the titled compound (231 mg). Melting point: 160-163 °C (crystallization solvent: ethyl acetate - n-hexane) FAB(pos) 426.3 [M+H]+

Example 204

N-[6-[2-(1-Pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



- 5 Concentrated hydrochloric acid (2 ml) was added to N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl]acetamide (98.0 mg, 0.345 mmol) obtained in Reference Example 103, which was stirred at 100 °C for 16 hours. The solvent was distilled out under reduced
- 10 pressure. Ethyl acetate was added to the residue, which was washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. WSC (62.5 mg, 0.326
- 15 mmol) was added to a dimethylformamide solution (1.5ml) of the resulting oily substance (79.0 mg, 0.326 mmol), [1,1'-biphenyl]-4-carboxylic acid (64.6 mg, 0.326 mmol) and DMAP (39.8 mg, 0.326 mmol) under ice-cooling, which was stirred at room temperature for 1 day. Ethyl acetate was
- 20 added to the reaction mixture, washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, then the solvent was distilled out under reduced pressure. The resulting residue was purified by aluminum column
- 25 chromatography (development solvent; ethyl acetate), powdered with ethyl acetate and isopropyl ether (1:5), to give the titled compound (36.8 mg).
- ¹H NMR (DMSO-d₆) δ: 1.67 (4H, m), 2.23 (2H, m), 2.34 (2H, m), 2.46 (4H, m), 2.57 (2H, m), 2.75 (2H, m), 6.24 (1H, s), 6.98 (1H, d, J = 8.1 Hz), 7.40-7.59 (5H, m), 7.76 (2H, d, J = 7.5 Hz), 7.82 (2H, d, J=8.4 Hz), 8.05 (2H, d, J = 8.4
- 30

WO 01/21577

PCT/JP00/06375

284

Hz), 10.19 (1H, s).


Melting point: 184-186 °C (crystallization solvent: ethyl acetate - isopropyl ether)

FAB(pos). 423.2 [M+H]⁺

5

Example 205

4'-Fluoro-N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



- 10 Concentrated hydrochloric acid (2 ml) was added to N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl]acetamide (98.0 mg, 0.345 mmol) obtained in Reference Example 103, which was stirred at 100°C for 16 hours. The solvent was distilled out under reduced
- 15 pressure. Ethyl acetate was added to the residue, which was washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. WSC (62.5 mg, 0.326
- 20 mmol) was added to a dimethylformamide solution (1.5 ml) of the resulting oily substance (79.0 mg, 0.326 mmol), 4'-fluoro-[1,1'-biphenyl]-4-carboxylic acid (64.6 mg, 0.326 mmol) and DMAP (39.8 mg, 0.326 mmol) under ice-cooling, which was stirred at room temperature for 1 day.
- 25 Ethyl acetate was added to the reaction mixture, which was washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and the solvent was distilled out under reduced pressure. The resulting residue was
- 30 purified by aluminum column chromatography (development solvent; ethyl acetate), powdered with ethyl acetate -

WO 01/21577

PCT/JP00/06375

285

isopropyl ether (1:5) , to give the titled compound (75.1 mg).

¹H NMR (DMSO-d₆) δ: 1.68 (4H, m), 2.23 (2H, m), 2.35 (2H, m), 2.50 (4H, m), 2.59 (2H, m), 2.75 (2H, m), 6.24 (1H, s),
5 6.98 (1H, d, J = 8.1 Hz), 7.34 (2H, m), 7.56 (2H, m), 7.81 (4H, m), 8.04 (2H, d, J = 8.4 Hz), 10.19 (1H, s).


Melting point: 187-189°C (crystallization solvent: ethyl acetate - isopropyl ether)

FAB (pos) 441.3 [M+H]⁺

10

Example 206

4'-Chloro-N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



15

Concentrated hydrochloric acid (2 ml) was added to N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-naphthalenyl]acetamide (98.0 mg, 0.345 mmol) obtained in Reference Example 103, which was stirred at 100°C for 16 hours. The solvent was distilled out under reduced

20

pressure. Ethyl acetate was added to the residue, which was washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. WSC (62.5 mg, 0.326 mmol) was added to a dimethylformamide solution (1.5 ml) of the resulting oily substance (79.0 mg, 0.326 mmol), 4'-chloro-[1,1'-biphenyl]-4-carboxylic acid (64.6 mg, 0.326 mmol) and DMAP (39.8 mg, 0.326 mmol) under ice-cooling, which was stirred at room temperature for 1 day.

30

Ethyl acetate was added to the reaction mixture, which was washed with aqueous potassium carbonate solution and

WO 01/21577

PCT/JP00/06375

286

saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under reduced pressure. The resulting residue was purified by aluminum column chromatography (development solvent; ethyl acetate), powdered with ethyl acetate - isopropyl ether (1:5), to give the titled compound (78.4 mg).


¹H NMR (DMSO-d₆) δ: 1.67 (4H, m), 2.23 (2H, m), 2.34 (2H, m), 2.45 (4H, m), 2.57 (2H, m), 2.75 (2H, m), 6.24 (1H, s), 6.98 (1H, d, J = 8.1 Hz), 7.55 (4H, m), 7.80 (2H, d, J=8.4 Hz), 7.84 (2H, d, J=8.4 Hz), 8.05 (2H, d, J = 8.7 Hz), 10.20 (1H, s).

Melting point: 207-209°C (crystallization solvent: ethyl acetate - isopropyl ether)

FAB (pos) 457.2 [M+H]⁺

Example 207

4'-Cyano-N-[6-[(dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Example 1, using N-[(6-amino-1,2,3,4-tetrahydro-2-naphthalenyl)methyl]-N,N-dimethylamine and 4'-cyano-[1,1'-biphenyl]-4-carboxylic acid.

¹H NMR (CDCl₃) δ: 1.42 (1H, m), 1.95 (2H, m), 2.26 (6H, s), 2.24-2.46 (3H, m), 2.84-2.95 (3H, m), 7.10 (1H, d, J=8.4 Hz), 7.30 (1H, m), 7.46 (1H, s), 7.74 (7H, m), 7.98 (2H, d, J=8.4 Hz).

Melting point: 183-185°C (crystallization solvent: ethyl acetate - isopropyl ether)

FAB (pos) 410.2 [M+H]⁺


WO 01/21577

PCT/JP00/06375

287

Example 208

N-[6-[2-(Dimethylamino)ethyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



5

Concentrated hydrochloric acid (1.5 ml) was added to N-[6-[2-(dimethylamino)ethyl]-7,8-dihydro-2-naphthalenyl]acetamide (57.5 mg, 0.223 mmol) obtained in Reference Example 104, which was stirred at 100°C for 1
10 hour. The solvent was distilled out under reduced pressure. Ethyl acetate was added to the residue, which was washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was
15 distilled out under reduced pressure. WSC (29.2 mg, 0.139 mmol) was added to a dimethylformamide solution (0.7 ml) of the resulting oily substance (30 mg, 0.139 mmol), [1,1'-biphenyl]-4-carboxylic acid (30.2 mg, 0.139 mmol) and DMAP (16.9 mg, 0.139 mmol) under ice-cooling, which was
20 stirred at room temperature for 16 hours. Ethyl acetate was added to the reaction mixture, which was washed with aqueous potassium carbonate solution and saturated aqueous sodium chloride solution, dried over anhydrous sodium sulfate, and then the solvent was distilled out under
25 reduced pressure. The resulting residue was purified by aluminum column chromatography (development solvent; ethyl acetate), powdered with ethyl acetate - isopropyl ether (1:5), to give the titled compound (12.4 mg).
¹H NMR (CDCl₃) δ: 2.29 (8H, m), 2.41 (2H, m), 2.46 (2H, m),
30 2.84 (2H, t, J = 8.1 Hz), 6.24 (1H, s), 6.98 (1H, d, J = 8.4 Hz), 7.34 (1H, m), 7.41 (1H, d, J = 6.9 Hz), 7.46 (3H,

WO 01/21577

PCT/JP00/06375

288


m), 7.63 (2H, d, J = 7.2 Hz), 7.71 (2H, d, J = 8.4 Hz), 7.77 (1H, br), 7.94 (2H, d, J = 8.4 Hz).

Melting point: 148-150°C (crystallization solvent: ethyl acetate - isopropyl ether)

5 FAB (pos) 397.2 [M+H]⁺

Example 209

N-[6-[2-(Dimethylamino)ethyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



10

A methanol solution (5 ml) of N-[6-[2-(dimethylamino)ethyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide (20 mg, 0.050 mmol) obtained in Example 208 and palladium carbon (10 mg) was stirred under hydrogen atmosphere for 4 hours. After a catalyst was filtered off, the filtrate was concentrated under reduced pressure. The resulting residue was purified by aluminum B column chromatography (development solvent; ethyl acetate), powdered with ethyl acetate - hexane (1:3), to give the titled compound (4.0 mg).

15

¹H NMR (CDCl₃) δ: 1.60 (4H, m), 1.92 (1H, m), 2.26 (6H, s), 2.42 (3H, m), 2.84 (3H, m), 7.06 (1H, d, J=8.1Hz), 7.32 (1H, m), 7.46 (4H, m), 7.63 (2H, d, J=6.9Hz), 7.72 (3H, m), 7.94 (2H, d, J=8.1Hz).

20

Melting point: 112-114°C (crystallization solvent: ethyl acetate - isopropyl ether)

25

FAB(pos) 399.2 [M+H]⁺

Example 210


4'-Chloro-N-[2-[(dimethylamino)methyl]-3,4-dihydro-2H-1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide

30

WO 01/21577

PCT/JP00/06375

289




The titled compound was obtained as white powders by the same method as in Example 1, using 6-amino-2-(dimethylamino)methyl-1,4-benzoxazin obtained in Reference Example 105.

¹H-NMR (CDCl₃) δ: 2.33 (6H, s), 2.44-2.65 (2H, m), 3.15-3.21 (1H, m), 3.41-3.46 (1H, m), 3.87 (1H, brs), 4.24-4.26 (1H, m), 6.61 (1H, dd, J=2.5, 8.6 Hz), 6.81 (1H, d, J=8.6 Hz), 7.28 (1H, d, J=2.5 Hz), 7.43 (2H, d, J=6.5 Hz), 7.54 (2H, d, J=6.5 Hz), 7.64 (2H, d, J=8.4 Hz), 7.71 (1H, s), 7.90 (2H, d, J=8.4 Hz).

Melting point: 227-230 °C (crystallization solvent: diisopropyl ether)

Example 211

4'-Methoxy-N-[6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 106.

¹H NMR (CDCl₃) δ: 2.31 (3H, s), 2.33 (2H, t, J=8.1 Hz), 2.49 (8H, bs), 2.84 (2H, t, J=8.1 Hz), 3.07 (2H, s), 3.87 (3H, s), 6.36 (1H, s), 7.00-7.03 (3H, m), 7.36 (1H, d, J=8.1 Hz), 7.51 (1H, s), 7.58 (2H, d, J=8.4 Hz), 7.67 (2H, d, J=8.4 Hz), 7.78 (1H, s), 7.91 (2H, d, J=8.4 Hz).

WO 01/21577


PCT/JP00/06375

290

Melting point: 208-210 °C (crystallization solvent : ethyl acetate)

Example 212

5 6-(4-Methoxyphenyl)-N-[6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]nicotinamide



10 The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 106.


¹H NMR (CDCl₃) δ : 2.30 (3H, s), 2.33 (2H, t, J=8.1 Hz), 2.47 (8H, bs), 2.84 (2H, t, J=8.1 Hz), 3.07 (2H, s), 3.89 (3H, s), 6.36 (1H, s), 7.01-7.04 (3H, m), 7.37 (1H, d, J=8.1 Hz), 7.49 (1H, s), 7.78-7.81 (2H, m), 8.03 (2H, d, J=8.4 Hz), 8.21 (1H, dd, J=2.1 Hz, 8.7 Hz), 9.09 (1H, s).

Melting point: 235-237 °C (crystallization solvent : ethyl acetate)

20

Example 213

N-[4-Methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide



25 The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using

WO 01/21577

PCT/JP00/06375

291


4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine
obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.77 (4H, s), 2.05 (3H, s), 2.51 (4H, s),
3.25 (2H, s), 4.74 (2H, s), 7.14-7.50 (6H, m), 7.63 (2H,
5 d, J=7.2 Hz), 7.71 (2H, d, J=8.4 Hz), 7.79 (1H, s), 7.94
(2H, d, J=8.4 Hz).

Melting point: 176-178 °C (crystallization solvent:
ethyl acetate - diisopropyl ether)

10 Example 214

4'-Methoxy-N-[4-methyl-3-(1-pyrrolidinylmethyl)-2H-
chromen-7-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as colorless powders
15 by carrying out the same operation as in Example 1, using
4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine
obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.77 (4H, s), 2.05 (3H, s), 2.51 (4H, s),
3.25 (2H, s), 3.87 (3H, s), 4.74 (2H, s), 7.01 (2H, d, J=8.7
20 Hz), 7.14-7.31 (3H, m), 7.57 (2H, d, J=8.7 Hz), 7.66 (2H,
d, J=8.4 Hz), 7.89 (1H, s), 7.91 (2H, d, J=8.4 Hz).

Melting point: 195-197 °C (crystallization solvent:
ethyl acetate - diisopropyl ether)


25 Example 215

N-[4-Methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl]-
6-phenylnicotinamide

WO 01/21577

PCT/JP00/06375

292



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine

5 obtained in Reference Example 107.


$^1\text{H NMR}$ (CDCl_3) δ : 1.77 (4H, s), 2.05 (3H, s), 2.51 (4H, s), 3.25 (2H, s), 4.74 (2H, s), 7.14-7.28 (3H, m), 7.47-7.54 (3H, m), 7.81-7.87 (2H, m), 8.06 (2H, d, $J=8.4$ Hz), 8.27 (1H, d, $J=8.4$ Hz), 9.13 (1H, s).

10 Melting point: 192-193 °C (crystallization solvent: ethyl acetate)

Example 216

6-(4-Methoxyphenyl)-N-[4-methyl-3-(1-

15 pyrrolidinylmethyl)-2H-chromen-7-yl]nicotinamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine

20 obtained in Reference Example 107.

$^1\text{H NMR}$ (CDCl_3) δ : 1.77 (4H, s), 2.05 (3H, s), 2.51 (4H, s), 3.25 (2H, s), 3.89 (3H, s), 4.74 (2H, s), 7.03 (2H, d, $J=8.7$ Hz), 7.14-7.26 (3H, m), 7.75-7.81 (2H, m), 8.03 (2H, d, $J=8.7$ Hz), 8.21 (1H, d, $J=6.6$ Hz), 9.09 (1H, s).

25 Melting point: 201-203 °C (crystallization solvent: ethyl

WO 01/21577


PCT/JP00/06375

293

acetate)

Example 217

N-[4-Methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl]-
5 4-phenyl-1-piperidinecarboxamide




The titled compound was obtained as colorless powders
by carrying out the same operation as in Example 99, using
4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine
10 obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.72-1.95 (8H, m), 2.03 (3H, s), 2.54 (4H,
s), 2.63-2.76 (1H, m), 2.95-3.00 (2H, m), 3.27 (2H, s),
4.19-4.23 (2H, m), 4.70 (2H, s), 6.39 (1H, s), 6.83 (1H,
s), 7.01-7.32 (7H, m).

15 Melting point: 125-127 °C (crystallization solvent:
ethyl acetate - diisopropyl ether)

Example 218

4-(4-Methoxyphenyl)-N-[4-methyl-3-(1-
20 pyrrolidinylmethyl)-2H-chromen-7-yl]-1-
piperidinecarboxamide



The titled compound was obtained as colorless powders
by carrying out the same operation as in Example 99, using
25 4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine

WO 01/21577

PCT/JP00/06375


294

obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.63-1.91 (8H, m), 2.02 (3H, s), 2.49 (4H, s), 2.61-2.71 (1H, m), 2.93-3.01 (2H, m), 3.23 (2H, s), 3.79 (3H, s), 4.16-4.21 (2H, m), 4.69 (2H, s), 6.34 (1H, s),
5 6.82-6.91 (3H, m), 6.99-7.02 (1H, m), 7.10-7.15 (3H, m).
Melting point: 144-146 °C (crystallization solvent: ethyl acetate - n-hexane)

Example 219

10 N-[4-Methyl-3-(4-morpholinylmethyl)-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using
15 4-methyl-3-(4-morpholinylmethyl)-2H-chromen-7-amine
obtained in Reference Example 108.

¹H NMR (DMSO-d₆) δ: 2.01 (3H, s), 2.37 (4H, s), 3.32 (2H, s), 3.57 (4H, s), 4.63 (2H, s), 7.23 (1H, d, J=8.1 Hz),
7.38-7.54 (5H, m), 7.76 (2H, d, J=7.5 Hz), 7.84 (2H, d, J=8.1
20 Hz), 8.04 (2H, d, J=8.1 Hz), 10.27 (1H, s).
Melting point: 162-164 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


Example 220

25 4'-Methoxy-N-[4-methyl-3-(4-morpholinylmethyl)-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide

WO 01/21577

PCT/JP00/06375

295




The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 4-methyl-3-(4-morpholinylmethyl)-2H-chromen-7-amine obtained in Reference Example 108.

¹H NMR (DMSO-d₆) δ: 2.00 (3H, s), 2.37 (4H, s), 3.11 (2H, s), 3.57 (4H, s), 3.82 (3H, s), 4.63 (2H, s), 7.07 (2H, d, J=8.7 Hz), 7.23 (1H, d, J=8.1 Hz), 7.38-7.40 (2H, m), 7.72 (2H, d, J=8.7 Hz), 7.79 (2H, d, J=8.4 Hz), 8.01 (2H, d, J=8.4 Hz), 10.23 (1H, s).

Melting point: 198-200 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 221

N-[6-(4-Morpholinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-(4-morpholinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 109.

¹H-NMR (CDCl₃) δ: 2.34 (2H, t, J=8.4 Hz), 2.45 (4H, m), 2.85 (2H, t, J=8.4 Hz), 3.06 (2H, s), 3.73 (4H, t, J=4.7 Hz), 6.36 (1H, s), 7.02 (1H, d, J=8.1 Hz), 7.36-7.78 (10H, m), 7.93 (2H, d, J=8.1 Hz).

Melting point: 180-181 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

WO 01/21577


PCT/JP00/06375

296

ethyl acetate - diisopropyl ether)

Example 222

6-(4-Methylphenyl)-N-[6-(4-morpholinylmethyl)-7,8-
5 dihydro-2-naphthalenyl]nicotinamide




The titled compound was obtained as colorless powders
by carrying out the same operation as in Example 1, using
6-(4-morpholinylmethyl)-7,8-dihydro-2-naphthalenamine
10 obtained in Reference Example 109.

¹H-NMR (CDCl₃) δ: 2.39 (2H, t, J=8.4 Hz), 2.43 (7H, m), 2.85
(2H, t, J=8.4 Hz), 3.06 (2H, s), 3.73 (4H, t, J=4.5 Hz),
6.36 (1H, s), 7.03 (1H, d, J=8.1 Hz), 7.30-7.38 (3H, m),
7.50 (1H, s), 7.76 (1H, s), 7.84 (1H, d, J=8.1 Hz), 7.97
15 (2H, d, J=8.1 Hz), 8.24 (1H, dd, J=8.4, 2.3 Hz), 9.12 (1H,
s).

Melting point: 233-234 °C (crystallization solvent: ethyl
acetate - diisopropyl ether)

20 Example 223

4-(4-Methylphenyl)-N-[6-(4-morpholinylmethyl)-7,8-
dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained as colorless powders
25 by carrying out the same operation as in Example 99, using
6-(4-morpholinylmethyl)-7,8-dihydro-2-naphthalenamine
obtained in Reference Example 109.

WO 01/21577

PCT/JP00/06375


297

¹H-NMR (CDCl₃) δ: 1.65-1.75 (4H, m), 1.90 (2H, m), 2.27-2.43 (7H, m), 2.72 (1H, m), 2.79 (2H, t, J=7.5 Hz), 2.93-3.04 (4H, m), 3.72 (4H, m), 4.20 (2H, d, J=11.7 Hz), 6.31 (1H, s), 6.39 (1H, s), 6.92 (1H, d, J=8.1 Hz), 7.05-7.26 (6H, m).

Melting point: 231-214 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 224

10 4'-Methyl-N-[6-(4-morpholinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-(4-morpholinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 109.

¹H-NMR (CDCl₃) δ: 2.33 (2H, t, J=8.1 Hz), 2.42-2.44 (7H, m), 2.84 (2H, t, J=8.1 Hz), 3.06 (2H, s), 3.72 (4H, t, J=4.2 Hz), 6.36 (1H, s), 7.01 (1H, d, J=8.1 Hz), 7.25-7.29 (2H, m), 7.37 (1H, d, J=8.1 Hz), 7.51-7.54 (3H, m), 7.68 (2H, d, J=8.1 Hz), 7.85 (1H, s), 7.92 (2H, d, J=8.1 Hz).

Melting point: 196-197 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


25 Example 225

2'-Methyl-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

WO 01/21577

PCT/JP00/06375


298



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
5 obtained in Reference Example 54.
Melting point: 177-178 °C (crystallization solvent : ethyl acetate - diisopropyl ether)

Example 226

10 4'-Fluoro-N-methyl-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide Hydrochloride



N-Methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine dihydrochloride (315 mg, 1.0 mmol)
15 obtained in Reference Example 113 was dissolved in N,N-dimethylformamide (25 ml). 4-Bromobenzoic acid (402 mg, 2.0 mmol), WSC (383 mg, 2.0 mmol), HOBT (270 mg, 2.0 mmol) and DMAP (244mg, 2.0 mmol) were added to the solution,
20 which was stirred at room temperature for 16 hours. Ethyl acetate and water were added to the reaction mixture, and extraction was conducted. The ethyl acetate layer was concentrated under reduced pressure. The residue was purified by aluminum column chromatography (development
25 solvent ; ethyl acetate : n-hexane = 33:67). The eluate was concentrated under reduced pressure, which was dissolved in dimethoxyethane - tetrahydrofuran (10:1, 5.5 ml).


WO 01/21577

PCT/JP00/06375

299

4-Fluorophenylboric acid (73 mg, 0.52 mmol) ,
tetrakis(triphenylphosphine)palladium complex (15 mg,
0.013 mmol) and 2N aqueous sodium carbonate solution (0.433 ml) were added to the solution, which was refluxed
5 with heating under nitrogen atmosphere at 90°C for 5.5
hours. The reaction mixture was poured into cold water,
and extraction was conducted using ethyl acetate. The
ethyl acetate layer was concentrated, and the residue was
purified by aluminum column chromatography (development
10 solvent; ethyl acetate) . 4N Hydrogen chloride - ethyl
acetate solution was added to the eluate, which was
concentrated under reduced pressure. The resulting
residue was recrystallized from methanol - ethyl acetate,
to give the titled compound (108 mg) .
15 ¹H-NMR (DMSO-d₆) δ: 1.92-1.98 (4H, m), 2.39 (2H, t, J=8.1
Hz), 2.73 (2H, t, J=8.1 Hz), 3.00 (2H, m), 3.35 (3H, m),
3.44 (2H, m), 3.83 (2H, d, J=5.6 Hz), 6.62 (1H, s), 6.92-7.01
(2H, m), 7.11 (1H, s), 7.26 (2H, dd, J=8.9, 5.6 Hz), 7.38
(2H, d, J=8.1 Hz), 7.55 (2H, d, J=8.1 Hz), 7.69 (2H, dd,
20 J=8.9, 5.6 Hz), 10.60 (1H, brs.).
Melting point: 201-203 °C (crystallization solvent :
methanol - diisopropyl ether)
FAB(pos) 441.2 [M+H]⁺

25 Example 227
(E)-3-(4-Chlorophenyl)-N-[6-[(dimethylamino)methyl]-
5,6,7,8-tetrahydro-2-naphthalenyl]-2-propenamide
Hydrochloride



30 The titled compound was obtained as colorless powders
by carrying out the same operation as in Example 4.
Melting point: 243-245 °C (crystallization solvent :
methanol - diisopropyl ether)


WO 01/21577

PCT/JP00/06375

300

Example 228


6-(4-Methylphenyl)-N-[5-methyl-6-(1-
pyrrolidinylmethyl)-7,8-dihydro-2-
5 naphthalenyl]nicotinamide



The titled compound was obtained as colorless powders
by carrying out the same operation as in Example 1, using
5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-
10 naphthalenamine obtained in Reference Example 69.
Melting point: 175-176 °C (crystallization solvent: ethyl
acetate - diisopropyl ether)
Elemental analysis for C₂₉H₃₀N₃O
Calcd.: C, 79.78; H, 6.93; N, 9.63
15 Found: C, 79.66; H, 6.97; N, 9.68

Example 229

4'-Fluoro-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-
dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as colorless powders
by carrying out the same operation as in Example 1, using
5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-
naphthalenamine obtained in Reference Example 69.
25 Melting point: 199-201 °C (crystallization solvent: ethyl

WO 01/21577

PCT/JP00/06375

301

acetate - diisopropyl ether)

Elemental analysis for $C_{29}H_{30}FN_2O$


Calcd.: C, 79.06; H, 6.63; N, 6.36

Found: C, 79.01; H, 6.81; N, 6.45

5

Example 230

6-(4-Fluorophenyl)-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]nicotinamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 69.

15 Melting point: 204-205 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Elemental analysis for $C_{28}H_{28}FN_3O$


Calcd.: C, 76.17; H, 6.39; N, 9.52

Found: C, 76.03; H, 6.44; N, 9.62

20

Example 231

4-(4-Fluorophenyl)-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



WO 01/21577

PCT/JP00/06375


302

The titled compound was obtained as colorless powders by carrying out the same operation as in Example 99, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 69.

5 Melting point: 172-173 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 232

4'-Methyl-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide




The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

15 Melting point: 176-177 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 233

20 N-[5-Methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-6-phenylnicotinamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 69.

25

WO 01/21577

PCT/JP00/06375

303

Melting point: 178-179 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


Elemental analysis for C₂₈H₂₉N₃O

Calcd.: C, 79.40; H, 6.90; N, 9.92

5 Found: C, 79.13; H, 6.82; N, 10.03

Example 234

4'-Methoxy-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 69.

15 ¹H-NMR (CDCl₃) δ: 1.78 (4H, m), 2.10 (3H, s), 2.37 (2H, t, J=8.1 Hz), 2.53 (4H, m), 2.76 (2H, t, J=8.1 Hz), 3.28 (2H, s), 3.87 (3H, s), 7.01 (1H, d, J=8.6 Hz), 7.27 (2H, d, J=7.8 Hz), 7.46 (1H, d, J=7.8 Hz), 7.48 (1H, s), 7.57 (2H, d, J=8.6 Hz), 7.66 (2H, d, J=8.6 Hz), 7.81 (1H, s), 7.92 (2H, d, J=7.8 Hz).

Melting point: 179-180 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Elemental analysis for C₃₀H₃₂N₂O₂

Calcd.: C, 79.61; H, 7.13; N, 6.19

25 Found: C, 79.35; H, 7.28; N, 6.24

Example 235


4-(4-Methoxyphenyl)-N-[5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide

30

WO 01/21577

PCT/JP00/06375

304



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 99, using 5-methyl-6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-

5 naphthalenamine obtained in Reference Example 69.

¹H-NMR (CDCl₃) δ: 1.67 (2H, dd, J=13.4, 4.0 Hz), 1.78 (4H, m), 1.89 (2H, d, J=11.4 Hz), 2.07 (3H, s), 2.34 (2H, t, J=7.5 Hz), 2.52 (4H, m), 2.68-2.73 (3H, m), 2.98 (2H, t, J=7.5 Hz), 3.26 (2H, s), 3.80 (3H, s), 4.20 (2H, d, J=13.4 Hz), 6.36 (1H, s), 6.86 (2H, d, J=8.4 Hz), 7.12-7.20 (5H, m).

Melting point: 163-164 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


Elemental analysis for C₂₈H₃₇N₃O₂

Calcd.: C, 75.13; H, 8.33; N, 9.39

15 Found: C, 74.96; H, 8.14; N, 9.10

Example 236

4-(4-Methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



20

The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

25 ¹H-NMR (CDCl₃) δ: 1.61-1.91 (8H, m), 2.31 (2H, t, J=8.1 Hz), 2.54 (4H, m), 2.73-2.81 (3H, m), 2.98 (2H, t, J=7.8 Hz),

WO 01/21577

PCT/JP00/06375


305

3.16 (2H, s), 3.79 (3H, s), 4.20 (2H, d, J=13.1 Hz), 6.31 (1H, s), 6.36 (1H, s), 6.86 (2H, d, J=8.6 Hz), 7.06-7.20 (5H, m).

Melting point: 175-176 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 237

4-(Benzyloxy)-N-[6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]benzamide



The titled compound was obtained as colorless powders by carrying out the same operation as in Example 1, using 6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenamine obtained in Reference Example 54.

Melting point: 174-175 °C (crystallization solvent: ethyl acetate - diisopropyl ether)


Elemental analysis for $C_{28}H_{30}N_2O_2$

Calcd.: C, 78.84; H, 7.09; N, 6.87

Found: C, 79.06; H, 6.99; N, 6.41

Example 238

4-(4-Methylphenyl)-N-[5-methyl-6-(1-pyrrolidinymethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 5-methyl-6-(1-

WO 01/21577

PCT/JP00/06375

306

pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenamine
obtained in Reference Example 69.

¹H-NMR (CDCl₃) δ: 1.65-1.78 (6H, m), 1.90 (2H, d, J=12.9 Hz), 2.07 (3H, s), 2.33-2.37 (5H, m), 2.53 (4H, m),
5 2.68-2.74 (3H, m), 2.99 (2H, m), 3.27(2H,s), 4.21 (2H, d, J=13.2 Hz), 6.37 (1H, s), 7.09-7.21 (7H, m).


Melting point: 159-160 °C (crystallization solvent:
ethyl acetate - diisopropyl ether)

FAB(pos) 444.3 [M+H]⁺

10

Example 239

4-(4-Fluorophenyl)-N-[6-(1-piperidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



15 The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-piperidinylmethyl)-7,8-dihydro-2-naphthalenamine dihydrochloride obtained in Reference Example 114.

¹H-NMR (CDCl₃) δ: 1.43 (2H, m), 1.56-1.75 (6H, m), 1.89 (2H, d, J=12.3 Hz), 2.27-2.36 (6H, m), 2.70 (1H, m), 2.78 (2H, t, J=7.5 Hz), 2.88-3.00 (4H, m), 4.20 (2H, d, J=13.2 Hz),
20 6.29 (1H, s), 6.38 (1H, s), 6.91-7.08 (4H, m), 7.14-7.20(3H, m).

Melting point: 194 -195 °C (crystallization solvent:
25 ethyl acetate - diisopropyl ether)


Example 240

4-(4-Methylphenyl)-N-[6-(1-piperidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide

WO 01/21577

PCT/JP00/06375

307



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-(1-piperidinylmethyl)-7,8-dihydro-2-naphthalenamine

5 dihydrochloride obtained in Reference Example 114.


¹H-NMR (CDCl₃) δ: 1.43 (2H, m), 1.56-1.74 (6H, m), 1.90 (2H, d, J=12.0 Hz), 2.27-2.36 (9H, m), 2.69 (1H, m), 2.79 (2H, t, J=8.1 Hz), 2.94-3.01 (4H, m), 4.19 (2H, d, J=13.2 Hz), 6.29 (1H, s), 6.35 (1H, s), 6.93 (2H, d, J=8.1 Hz), 7.05-7.26 (5H, m).

10

Melting point: 209 -210 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

Example 241

15 4-(4-Methylphenyl)-N-[6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine

20

obtained in Reference Example 106.
¹H NMR (CDCl₃) δ: 1.62-1.77 (2H, m), 1.90 (2H, d, J=12.0 Hz), 2.28 (2H, t, J=8.1 Hz), 2.29 (3H, s), 2.33 (3H, s), 2.46 (8H, bs), 2.64-2.73 (1H, m), 2.79 (2H, t, J=8.1 Hz), 2.96 (2H, d, J=10.5 Hz), 3.05 (2H, s), 4.19 (2H, d, J=13.5 Hz), 6.31 (1H, s), 6.34 (1H, s), 6.93 (1H, d, J=8.4 Hz), 7.04-7.16 (5H, m), 7.23 (1H, s).

25

WO 01/21577

PCT/JP00/06375

308

Melting point: 214-216 °C (crystallization solvent:
tetrahydrofuran - n-hexane)


Elemental analysis for C₂₉H₃₈N₄O

Calcd.: C, 75.94; H, 8.35; N, 12.22.

5 Found: C, 75.67; H, 8.47; N, 12.27.

Example 242

4-(4-Methoxyphenyl)-N-[6-[(4-methyl-1-
piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-1-
10 piperidinecarboxamide



The titled compound was obtained by carrying out the
same operation as in Example 99, using 6-[(4-methyl-1-
piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine
15 obtained in Reference Example 106.

¹H NMR (CDCl₃) δ: 1.68-1.76 (2H, m), 1.89 (2H, d, J=11.1
Hz), 2.29 (2H, t, J=8.1 Hz), 2.29 (3H, s), 2.46 (8H, bs),
2.64-2.71 (1H, m), 2.79 (2H, t, J=8.1 Hz), 2.82-3.03 (2H,
m), 3.05 (2H, s), 3.80 (3H, s), 4.19 (2H, d, J=12.6 Hz),
20 6.31 (1H, s), 6.34 (1H, s), 6.87 (2H, d, J=8.7 Hz), 6.93
(1H, d, J=8.4 Hz), 7.06 (1H, dd, J=8.1, 2.1 Hz), 7.14 (2H,
d, J=8.7 Hz), 7.23 (1H, s).

Melting point: 198-200 °C (crystallization solvent:
tetrahydrofuran - n-hexane)

25 Elemental analysis for C₂₉H₃₈N₄O₂

Calcd.: C, 73.38; H, 8.07; N, 11.80.

Found: C, 73.04; H, 7.95; N, 11.67.


Example 243

30 4-(4-Chlorophenyl)-N-[6-[(4-methyl-1-
piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-1-
piperidinecarboxamide

WO 01/21577

PCT/JP00/06375

309



The titled compound was obtained by carrying out the same operation as in Example 99, using 6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine
5 obtained in Reference Example 106.


¹H NMR (CDCl₃) δ: 1.64-1.76 (2H, m), 1.90 (2H, d, J=11.1 Hz), 2.29 (2H, t, J=8.1 Hz), 2.29 (3H, s), 2.46 (8H, bs), 2.66-2.72 (1H, m), 2.79 (2H, t, J=8.1 Hz), 2.81-3.03 (2H, m), 3.05 (2H, s), 4.20 (2H, d, J=12.6 Hz), 6.31 (1H, s),
10 6.34 (1H, s), 6.93 (1H, d, J=7.8 Hz), 7.04-7.07 (1H, m), 7.14 (2H, d, J=8.4 Hz), 7.22 (1H, s), 7.28 (2H, d, J=8.4 Hz).

Melting point: 201-203 °C (crystallization solvent: tetrahydrofuran - n-hexane)

15

Example 244

N-[2-[(Dimethylamino)methyl]-1H-inden-6-yl][1,1'-biphenyl]-4-carboxamide



20

The titled compound was obtained by carrying out the same operation as in Example 1, using 2-[(dimethylamino)methyl]-1H-inden-6-amine obtained in Reference Example 116.

Elemental analysis for C₂₅H₂₄N₂O · 0.5H₂O

25

Calcd.: C, 79.55; H, 6.68; N, 7.42.

Found: C, 79.38; H, 6.76; N, 7.34.

Melting point: 187-189 °C (crystallization solvent: ethyl acetate - diisopropyl ether)

WO 01/21577


PCT/JP00/06375

310

FAB(pos) 369.2 [M+H]⁺

Example 245

N-[2-[(Dimethylamino)methyl]-1H-inden-6-yl]-4'-
5 fluoro[1,1'-biphenyl]-4-carboxamide




The titled compound was obtained by carrying out the
same operation as in Example 1, using 2-
[(dimethylamino)methyl]-1H-inden-6-amine obtained in
10 Reference Example 116.

Melting point: 209-211 °C (crystallization solvent: ethyl
acetate - diisopropyl ether)

FAB(pos) 387.2 [M+H]⁺

15 Example 246

4'-Chloro-N-[2-[(dimethylamino)methyl]-1H-inden-6-
yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the
20 same operation as in Example 1, using 2-
[(dimethylamino)methyl]-1H-inden-6-amine obtained in
Reference Example 116.

Melting point: 218-220 °C (crystallization solvent: ethyl
acetate - diisopropyl ether)

25 FAB(pos) 403.2 [M+H]⁺

Example 247


4'-Chloro-N-[2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-

WO 01/21577

PCT/JP00/06375

311

1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide




The titled compound was obtained by carrying out the same operation as in Example 1, using 6-amino-2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-1,4-benzoxazine obtained in Reference Example 117.

¹H-NMR (CDCl₃) δ: 1.70-1.90 (4H, m), 2.50-2.70 (4H, m), 2.73 (2H, d, J=6.0Hz), 3.18-3.24 (1H, m), 3.45-3.49 (1H, m), 3.87 (1H, brs), 4.26-4.28 (1H, m), 6.61 (1H, dd, J=2.7, 8.4 Hz), 6.80 (1H, d, J=8.4 Hz), 7.26 (1H, d, J=2.7 Hz), 7.44 (2H, d, J=8.4 Hz), 7.55 (2H, d, J=8.4 Hz), 7.64 (2H, d, J=8.1 Hz), 7.71 (1H, s), 7.91 (2H, d, J=8.1 Hz).
Melting point: 221-222 °C (crystallization solvent: diisopropyl ether)

Example 248

4'-Fluoro-N-[2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 6-amino-2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-1,4-benzoxazine obtained in Reference Example 117.

¹H-NMR (CDCl₃) δ: 1.70-1.90 (4H, m), 2.50-2.70 (4H, m), 2.73 (2H, d, J = 6.3Hz), 3.18-3.24 (1H, m), 3.45-3.49 (1H, m), 3.88 (1H, brs), 4.24-4.30 (1H, m), 6.62 (1H, dd, J=2.7, 8.4 Hz), 6.80 (1H, d, J=8.4 Hz), 7.13-7.19 (2H, m), 7.26 (1H,

WO 01/21577

PCT/JP00/06375

312


d, J=2.7 Hz), 7.56-7.60 (2H, m), 7.63 (2H, d, J=8.4 Hz), 7.71 (1H, s), 7.90 (2H, d, J=8.4 Hz).

Melting point: 204-206 °C (crystallization solvent: diisopropyl ether)

5

Example 249

6-(4-Methylphenyl)-N-[2-(1-pyrrolidinymethyl)-3,4-dihydro-2H-1,4-benzoxazin-6-yl]nicotinamide



10

The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 6-amino-2-(1-pyrrolidinymethyl)-3,4-dihydro-2H-1,4-benzoxazine obtained in Reference Example 117.


¹H-NMR (CDCl₃) δ: 1.70-1.85 (4H, m), 2.43 (3H, s), 2.50-2.70 (4H, m), 2.74 (2H, d, J=6.3Hz), 3.19-3.25 (1H, m), 3.45-3.49 (1H, m), 3.90 (1H, brs), 4.27-4.29 (1H, m), 6.63 (1H, dd, J=2.4, 8.7 Hz), 6.81 (1H, d, J=8.7 Hz), 7.26 (1H, d, J=2.7 Hz), 7.31 (2H, d, J=8.1 Hz), 7.67 (1H, s), 7.81 (1H, d, J=8.1 Hz), 7.93 (2H, d, J=7.8Hz), 8.21 (1H, dd, J=2.4, 8.4 Hz), 9.09 (1H, d, J=2.4 Hz).

20

Melting point: 207-208 °C (crystallization solvent: diisopropyl ether)

Example 250

4-(4-Fluorophenyl)-N-[2-(1-pyrrolidinymethyl)-3,4-dihydro-2H-1,4-benzoxazin-6-yl]-1-piperidinecarboxamide



25

WO 01/21577

PCT/JP00/06375

313


The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 6-amino-2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-1,4-benzoxazine obtained in Reference Example 117.

5 ¹H-NMR (CDCl₃) δ: 1.60-1.90 (8H, m), 2.50-2.70 (5H, m), 2.71 (2H, d, J=6.3Hz), 2.91-3.00 (2H, m), 3.15-3.21 (1H, brs), 3.42-3.45 (1H, m), 3.77 (1H, brs), 4.15-4.25 (3H, m), 6.20 (1H, s), 6.38 (1H, dd, J=2.1, 8.4 Hz), 6.73 (1H, d, J=8.4 Hz), 6.91 (1H, d, J=2.1 Hz), 6.97-7.03 (2H, m), 7.14-7.19 (2H, m).

Melting point: 192-195 °C (crystallization solvent: diisopropyl ether)

Example 251

15 4'-Chloro-N-[4-(methylsulfonyl)-2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide



20 The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 6-amino-4-(methylsulfonyl)-2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-1,4-benzoxazine obtained in Reference Example 118.

25 ¹H-NMR (CDCl₃) δ: 1.75-1.85 (4H, m), 2.55-2.70 (4H, m), 2.78 (2H, d, J=6.0Hz), 3.04 (3H, s), 3.27-3.34 (1H, m), 4.24-4.31 (1H, m), 4.31-4.35 (1H, m), 6.98 (1H, d, J=8.7 Hz), 7.45 (2H, d, J=9.0 Hz), 7.50-7.60 (1H, m), 7.53 (2H, d, J=9.0 Hz), 7.67 (2H, d, J=8.4 Hz), 7.84 (1H, s), 7.84 (1H, brs), 7.94 (2H, d, J=8.4 Hz).

30 Melting point: 203-204 °C (crystallization solvent: diisopropyl ether)


WO 01/21577

PCT/JP00/06375

314

Example 252

N-[6-[(4-Methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide




5 The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 106.

¹H NMR (CDCl₃) δ: 2.31 (3H, s), 2.33 (2H, t, J=8.1 Hz), 2.49 (8H, bs), 2.84 (2H, t, J=8.1 Hz), 3.07 (2H, s), 6.36 (1H, s), 7.02 (1H, d, J=8.1 Hz), 7.35-7.52 (5H, m), 7.63 (2H, d, J=8.1 Hz), 7.71 (2H, d, J=8.1 Hz), 7.80 (1H, s), 7.94 (2H, d, J=8.1 Hz).

10 Melting point: 196-198 °C (crystallization solvent: ethyl acetate)

Example 253

4'-Methyl-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



25 The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

¹H NMR (CDCl₃) δ: 2.08 (3H, s), 2.29 (3H, s), 2.34 (2H, t, J=7.8 Hz), 2.42 (3H, s), 2.45 (8H, bs), 2.75 (2H, t, J=7.8 Hz), 3.16 (2H, s), 7.26-7.30 (3H, m), 7.44 (1H, d, J=8.4

WO 01/21577

PCT/JP00/06375

315


Hz), 7.53-7.55 (3H, m), 7.70 (2H, d, J=8.4 Hz), 8.00 (1H, s), 7.93 (2H, d, J=8.4 Hz).

Melting point: 212-214 °C (crystallization solvent: ethyl acetate)

5

Example 254

4'-Methoxy-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



10

The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

15 ¹H NMR (CDCl₃) δ: 2.08 (3H, s), 2.29 (3H, s), 2.34 (2H, t, J=7.8 Hz), 2.45 (8H, bs), 2.75 (2H, t, J=7.8 Hz), 3.16 (2H, s), 3.87 (3H, s), 7.01 (2H, d, J=8.1 Hz), 7.27 (1H, d, J=8.4 Hz), 7.44 (1H, d, J=8.4 Hz), 7.51 (1H, s), 7.58 (2H, d, J=8.4 Hz), 7.67 (2H, d, J=8.4 Hz), 7.81 (1H, s), 7.92 (2H, d, J=8.4 Hz).

20

Melting point: 215-217 °C (crystallization solvent: ethyl acetate)


Example 255

25 4'-Fluoro-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide

WO 01/21577

PCT/JP00/06375

316




The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

¹H NMR (CDCl₃) δ: 2.08 (3H, s), 2.29 (3H, s), 2.34 (2H, t, J=7.8 Hz), 2.46 (8H, bs), 2.75 (2H, t, J=7.8 Hz), 3.16 (2H, s), 7.17 (2H, d, J=8.4 Hz), 7.28 (1H, d, J=8.4 Hz), 7.44 (1H, d, J=8.4 Hz), 7.51 (1H, s), 7.57-7.62 (2H, m), 7.66 (2H, d, J=8.4 Hz), 7.82 (1H, s), 7.94 (2H, d, J=8.4 Hz). Melting point: 233-235 °C (crystallization solvent: ethyl acetate)

Example 256

4'-Chloro-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide



The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

¹H NMR (CDCl₃) δ: 2.08 (3H, s), 2.29 (3H, s), 2.34 (2H, t, J=7.8 Hz), 2.46 (8H, bs), 2.75 (2H, t, J=7.8 Hz), 3.16 (2H, s), 7.28 (1H, d, J=8.4 Hz), 7.43-7.47 (3H, m), 7.51 (1H, s), 7.56 (2H, d, J=8.4 Hz), 7.67 (2H, d, J=8.4 Hz), 7.80 (1H, s), 7.94 (2H, d, J=8.4 Hz).

WO 01/21577


PCT/JP00/06375

317

Melting point: 216-218 °C (crystallization solvent: ethyl acetate)

Example 257

5 6-(4-Chlorophenyl)-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]nicotinamide



10 The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.


¹H NMR (CDCl₃) δ: 2.09 (3H, s), 2.29 (3H, s), 2.35 (2H, t, J=8.1 Hz), 2.46 (8H, bs), 2.75 (2H, t, J=8.1 Hz), 3.16 (2H, s), 7.28 (1H, d, J=8.4 Hz), 7.43-7.50 (4H, m), 7.83 (2H, d, J=8.4 Hz), 8.01 (2H, d, J=8.4 Hz), 8.27 (1H, d, J=8.4 Hz), 9.13 (1H, s).

Melting point: 219-221 °C (crystallization solvent: ethyl acetate)

20

Example 258

5-(4-Chlorophenyl)-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-2-pyridinecarboxamide



25

The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 5-

WO 01/21577

PCT/JP00/06375

318


methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

¹H NMR (CDCl₃) δ: 2.09 (3H, s), 2.29 (3H, s), 2.35 (2H, t, J=8.1 Hz), 2.45 (8H, bs), 2.77 (2H, t, J=8.1 Hz), 3.16 (2H, s), 7.30 (1H, d; J=8.1 Hz), 7.49-7.63 (6H, m), 8.05 (1H, dd, J=2.4 Hz, 8.4 Hz), 8.36 (1H, d, J=8.1 Hz), 8.79 (1H, d, J=1.2 Hz), 9.97 (1H, s).

Melting point: 177-179 °C (crystallization solvent: ethyl acetate)

Example 259

N-[5-Methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-4-(4-methylphenyl)-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Reference Example 99, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

¹H NMR (CDCl₃) δ: 1.60-1.78 (4H, m), 2.05 (3H, s), 2.28 (3H, s), 2.29 (2H, t, J=8.1 Hz), 2.33 (3H, s), 2.46 (8H, bs), 2.65-2.72 (3H, m), 2.93-3.03 (2H, m), 3.13 (2H, s), 4.18-4.23 (2H, m), 6.40 (1H, s), 7.09-7.24 (7H, m).

Melting point: 176-178 °C (crystallization solvent: ethyl acetate-hexane)


Example 260

4-(4-Methoxyphenyl)-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide

WO 01/21577

PCT/JP00/06375

319




The titled compound was obtained by carrying out the same operation as in Reference Example 99, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

¹H NMR (CDCl₃) δ : 1.68-1.92 (4H, m), 2.05 (3H, s), 2.28 (3H, s), 2.29 (2H, t, J=8.1 Hz), 2.45 (8H, bs), 2.67-2.72 (3H, m), 2.95-3.02 (2H, m), 3.14 (2H, s), 3.80 (3H, s), 4.18-4.22 (2H, m), 6.36 (1H, s), 6.87 (2H, d, J=8.4 Hz), 7.12-7.21 (5H, m).

Melting point: 175-177 °C (crystallization solvent: ethyl acetate)

Example 261

4-(4-Chlorophenyl)-N-[5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-1-piperidinecarboxamide



The titled compound was obtained by carrying out the same operation as in Reference Example 99, using 5-methyl-6-[(4-methyl-1-piperazinyl)methyl]-7,8-dihydro-2-naphthalenamine obtained in Reference Example 115.

¹H NMR (CDCl₃) δ : 1.67-1.92 (4H, m), 2.05 (3H, s), 2.28 (3H, s), 2.29 (2H, t, J=8.1 Hz), 2.45 (8H, bs), 2.67-2.72 (3H, m), 2.95-3.02 (2H, m), 3.14 (2H, s), 4.18-4.23 (2H, m), 6.36 (1H, s), 7.13-7.30 (7H, m).

Melting point: 141-143 °C (crystallization solvent: ethyl acetate)

WO 01/21577


PCT/JP00/06375

320

ethyl acetate)

Example 262

4-[(4-Chlorophenyl)(phenyl)methyl]-N-[4-methyl-3-(1-
pyrrolidinylmethyl)-2H-chromen-7-yl]-1-
piperazinecarboxamide




The titled compound was obtained by carrying out the
same operation as in Reference Example 99, using 4-
methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine
obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.76 (4H, s), 2.01 (3H, s), 2.42 (4H, t,
J=5.1 Hz), 2.49 (4H, s), 3.22 (2H, s), 3.48 (4H, t, J=5.1
Hz), 4.24 (1H, s), 4.68 (2H, s), 6.23 (1H, s), 6.77 (1H,
s), 6.96 (1H, d, J=8.7 Hz), 7.09 (1H, d, J=8.7 Hz), 7.19-7.61
(9H, m).

Melting point: 104-106 °C (crystallization solvent:
ethyl acetate - n-hexane)

Example 263

N-(2,2-Diphenylethyl)-N'-[4-methyl-3-(1-
pyrrolidinylmethyl)-2H-chromen-7-yl]urea



The titled compound was obtained by carrying out the
same operation as in Reference Example 99, using 4-
methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine

WO 01/21577

PCT/JP00/06375

321


obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.76 (4H, s), 1.99 (3H, s), 2.49 (4H, s),
3.22 (2H, s), 3.83 (2H, t, J=7.8 Hz), 4.18 (1H, t, J=7.8
Hz), 4.66 (2H, s), 4.96 (1H, s), 6.48 (1H, s), 6.57 (1H,
5 s), 6.69 (1H, d, J=8.1 Hz), 6.98 (1H, d, J=8.1 Hz), 7.20-7.30
(10H, m).

Melting point: 166-168 °C (crystallization solvent:
ethyl acetate - n-hexane)

10 Example 264

N-[4-Methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl]-
3,4-dihydro-2(1H)-isoquinolinecarboxamide




The titled compound was obtained by carrying out the
15 same operation as in Reference Example 99, using 4-
methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine
obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.76 (4H, s), 2.02 (3H, s), 2.49 (4H, s),
2.92 (2H, t, J=6.0 Hz), 3.23 (2H, s), 3.71 (2H, t, J=6.0 Hz),
20 4.65 (2H, s), 4.68 (2H, s), 6.43 (1H, s), 6.86 (1H, d, J=1.8
Hz), 7.02-7.22 (6H, m).

Melting point: 135-137 °C (crystallization solvent:
ethyl acetate - n-hexane)

25 Example 265

N-[4-Methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl]-
4-(1-piperidiny)-1-piperidinecarboxamide



WO 01/21577

PCT/JP00/06375


322

The titled compound was obtained by carrying out the same operation as in Reference Example 99, using 4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine obtained in Reference Example 107.

- 5 ¹H NMR (CDCl₃) δ: 1.27-1.89 (14H, m), 2.02 (3H, s), 2.49-2.51 (9H, m), 2.83-2.90 (2H, m), 3.23 (2H, s), 4.08-4.12 (2H, m), 4.68 (2H, s), 6.31 (1H, s), 6.80 (1H, d, J=2.4 Hz), 6.98 (1H, dd, J=2.4 Hz, 8.4 Hz), 7.09 (1H, d, J=8.4 Hz).
- 10 Melting point: 98-100 °C (crystallization solvent: ethyl acetate - n-hexane)

Example 266

- 2-(4-Methyl-6-oxo-2-phenyl-1,6-dihydro-5-pyrimidinyl)-
15 N-[4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-yl]acetamide



The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine obtained in Reference Example 107.

- 20 ¹H NMR (CDCl₃) δ: 1.76 (4H, s), 1.98 (3H, s), 2.49 (4H, s), 2.61 (3H, s), 3.22 (2H, s), 3.65 (2H, s), 4.65 (2H, s), 6.86-7.00 (4H, m), 7.54 (3H, s), 8.01 (2H, s), 8.87 (1H, s).
- 25 Melting point: 255-257 °C (crystallization solvent: ethyl acetate - n-hexane)


Example 267

- Benzyl 2-[[4-methyl-3-(1-pyrrolidinylmethyl)-2H-
30 chromen-7-yl]amino]-2-oxoethylcarbamate

WO 01/21577

PCT/JP00/06375

323



The titled compound was obtained by carrying out the same operation as in Reference Example 1, using 4-methyl-3-(1-pyrrolidinylmethyl)-2H-chromen-7-amine

5 obtained in Reference Example 107.

¹H NMR (CDCl₃) δ: 1.78 (4H, s), 2.03 (3H, s), 2.53 (4H, s), 3.26 (2H, s), 3.99 (2H, d, J=4.8 Hz), 4.71 (2H, s), 5.17 (2H, s), 5.50 (1H, bs), 7.00-7.14 (4H, m), 7.36 (5H, s), 7.80 (1H, bs).

10 Melting point: 143-145 °C (crystallization solvent: ethyl acetate - n-hexane)

Preparation Example 1

15	(1) Compound obtained in	
	Reference Example 25	50 mg
	(2) Lactose	34 mg
	(3) Corn starch	10.6 mg
	(4) Corn starch (paste)	5 mg
	(5) Magnesium stearate	0.4 mg
20	(6) <u>Carboxymethylcellulose calcium</u>	<u>20 mg</u>
	Total	120 mg

In accordance with a conventional manner, the above (1) to (6) are admixed and tableted using a tableting machine to give tablets.

25

Preparation Example 2

30	(1) Compound obtained in Example 1	50 mg
	(2) Lactose	34 mg
	(3) Corn starch	10.6 mg
	(4) Corn starch (paste)	5 mg
	(5) Magnesium stearate	0.4 mg
	(6) <u>Carboxymethylcellulose calcium</u>	<u>20 mg</u>
	Total	120 mg

In accordance with a conventional manner, the above (1) to (6) are admixed and tableted using a tableting machine to give tablets.

5 Reference Example 1-1

Amplification of rat SLC-1 receptor cDNA by PCR method using rat-brain-originated cDNA

Reverse transcription reaction was done using random primer, with rat-brain-originated poly (A)⁺RNA (Clone Tech
10 Co.) used as a template. Reagent from the TaKaRa RNA PCR ver. 2 kit was used for the reverse transcription reaction.

Next, using this reverse transcription product as a template, amplification was done by a PCR method using synthetic DNA primers with sequence numbers 1 and 2.

15 Synthetic DNA primer was constructed to amplify genes in the domain where genes are translated by receptor protein.

At that time, individual restriction enzyme recognition sequences were also added on the 5' side and 3' side of the gene, to add a nucleotide sequence on the 5' side of gene
20 which recognized restriction enzyme Sal I, and to add a nucleotide sequence on the 3' side of the gene which recognized the restriction enzyme Spe I. The reactant was constituted of 5 µl of cDNA template, 0.4 µM of synthetic DNA primer, 0.25 mM of dNTPs, 0.5 µl of Pfu (StrataGene Co.)
25 DNA polymerase, and buffers attached to enzymes, with total reaction quantity set at 50 µl.

A thermal cycler (Parkin Elmer Co.) was used to produce cycles for amplification. After heating at 94°C for 60 seconds, the cycle consisting of 94°C for 60 seconds, 60°C
30 for 30 seconds, and 72°C for 150 seconds, was repeated 35 times, and finally reaction was conducted at 72°C for 10 minutes. After 0.8% agarose gel electrophoresis, the amplified products were confirmed by ethidium bromide
35 dying.

Reference Example 1-2

Subcloning of PCR products into plasmid vector, and confirmation of an amplified cDNA sequence by decoding of a nucleotide sequence in an inserted cDNA portion

The reaction product after PCR conducted in Reference
5 Example 1-1 was separated using 0.8% low-melting point agarose gel. After the band section was cut out using a razor, DNA was recovered by conducting fragmentation, phenol extraction, phenol-chloroform extraction and ethanol precipitation. The recovered DNA was subcloned on
10 plasmid vector PCR-Script Amp SK(+) in accordance with prescription of the PCR-Script™ Amp SK(+) cloning kit (Stratagene Co.). After this was introduced into Escherichia coli XL-1 Blue (Stratagene Co.) by transformation, the clones with fragments of inserted cDNA
15 were selected in LB agar culture medium containing ampicillin and X-gal. Only clones showing white color were separated using a sterilized toothpick, and transformant E. coli XL-1 Blue/rat SLC-1 was obtained.

Each clone was cultured overnight in LB culture medium
20 containing ampicillin, and plasmid DNA was prepared using QIA prep8 mini prep (Qiagen). A portion of the prepared DNA was digested with Sal I and Spe I, and the size of the inserted receptor cDNA fragment was confirmed. Reactions to determine nucleotide sequences were carried out using
25 a DyeDeoxy Terminator Cycle Sequence Kit (Parkin Elmer Co.), and decoded using a fluorescent light automatic sequencer. The sequences of the 3 clones obtained were analyzed, and it was confirmed that all of them match the reported gene sequence (Sequence number: 4) in which the
30 Sal I recognition sequence is added on the 5' side and the Spe I recognition sequence is added on the 3' side of the cDNA sequence (Lakaye, B., et al., Biochim. Biophys. Acta, Vol. 1401, pp. 216-220 (1998), accession No. AF08650) coding rat SLC-1 protein (Sequence number: 3).

35

Reference Example 1-3

Preparation of CHO cells for rat SLC-1 expression

The full-length amino acid sequence of rat brain originated SLC-1, which was confirmed in Reference Example 1-2, was coded, and plasmid was prepared using a plasmid Midi Kit (Qiagen) from the E. coli transformed by the plasmid, to which the gene with Sal I recognition sequence added to the 5' side and Spe I recognition sequence added to the 3' side, had been introduced. Then, the insert section was cut out by digesting with Sal I and Spe I. The insert DNA was cut out with a razor from the agarose gel after electrophoresis.

Next, fragmentation, phenol extraction, phenol-chloroform extraction, and ethanol precipitation, were conducted and the DNA was recovered. This insert DNA was added to vector plasmid pAKKO-111H (the same vector plasmid as pAKKO1.11H described in Hinuma, S., et al., Biochim. Biophys. Acta, Vol. 1219, pp. 251-259 (1994)) for animal cell expression which was digested with Sal I and Spe I, and ligation was conducted using T4 ligase (TaKaRa Shuzo), to construct pAKKO-SLC-1 plasmid for protein expression.

After E. coli DH5 transformed by pAKKO-SLC-1 was cultured, pAKKO-SLC-1 plasmid DNA was prepared using a Plasmid Midi Kit (Qiagen). This was introduced into CHO dhfr⁻ cells in accordance with the attached protocol, using a CellPfect Transfection Kit (Amersham Pharmacia Biotech Co.). A coprecipitating suspension of 10 µg of DNA and calcium phosphate was prepared, and this suspension was added to 10 cm Petri dishes in which 5×10^5 or 1×10^6 of CHO dhfr⁻ cells had been seeded 24 hours previously. After these cells were cultured for 1 day in MEMa culture medium containing 10% fetal bovine serum, subculture was conducted, and cultivation was conducted in selective culture medium, MEMa culture medium containing no nucleic acid but containing 10% dialyzed fetal bovine serum. 56 clones of colonies of the transformed CHO cells expressing SLC-1, proliferated in the selective culture medium, were

selected.

Reference Example 1-4

Selection of CHO/SLC-1 cell strain expressing a large
5 quantity of full-length rat SLC-1 receptor protein mRNA

The quantity of expressed full-length rat SLC-1
receptor protein mRNA of 56 clones of the CHO/SLC-1 strains
established in Reference Example 1-3, was measured using
a Cytostar T Plate (Amersham Pharmacia Biotech Co.) as shown
10 below according to the attached protocol. Each well of the
Cytostar T Plate was seeded with each clone of the CHO/SLC-1
strain by 2.5×10^4 , and cultured for 24 hours, then the
cells were fixed using 10% formalin. After 0.25% Triton
X-100 was added to each well to increase cell permeability,
15 ^{35}S -labeled riboprobes with sequence number: 5 were added
and hybridized. 20 mg/ml of RNaseA was added to each well
to digest free riboprobes. After the plate was thoroughly
washed, the radioactivity of the hybridized riboprobes was
determined using a Topcounter. Strains with high
20 radioactivity showed large amounts of mRNA expression. In
particular, mainly used was Clone number 44 among 3 clones
which showed large amounts of mRNA expression.

Reference Example 1-5

25 Isolation of plasmid containing human SLC-1 cDNA

After nicks were inserted into the DNA of Human fetal
brain originated cDNA library (SUPERScript™ cDNA Library;
GIBCOBRL Co.) according to the manual of the Genetrapp
cDNA positive selection system (GIBCOBRL Co.), using phage
30 F1 endonuclease, single stranded human fetal brain
originated cDNA library was prepared by digesting the
above-mentioned library with Escherichia coli exonuclease
III.

Biotin-14-dCTP was added to the 3' end of synthetic
35 oligonucleotide (equivalent to 1434-1451 of accession No.
U71092), sequence number: 6 which was prepared according

to the report by Kolakowski Jr., et al. (Kolakowski Jr.,
et al. (1996) FEBS Lett. Vol. 398, pp. 253-258) using
Terminal Deoxynucleotidyl Transferase, and biotinlated
oligonucleotide was prepared. The above manual was
5 followed regarding composition of a reaction mixture and
reaction time.

After 4 µg of single stranded human fetal brain
originated cDNA library was kept at 95° C for 1 minute, the
library was rapidly cooled on ice. 20 ng of biotinlated
10 oligonucleotide was added, which was hybridized using the
attached hybridization buffer at 37° C. for 1 hour.
Streptoavidin beads were added to the mixture, then single
stranded human fetal brain originated cDNA hybridized by
biotinlated oligonucleotide, was isolated using a MAGNA-
15 SEP Magnetic Particle Separator (GIBCOBRL Co.). The
complementary strand was synthesized according to the
manual, using as primer 50 ng of synthetic oligonucleotide
(equivalent to 1011 - 1028 of accession No. U71092) of
sequence number: 7, prepared based on the report by
20 Kolakowski Jr., et al (Kolakowski Jr., et al. (1996) FEBS
Lett. Vol. 398, pp. 253-258), to give the double stranded
plasmid.

Reference Example 1-6

25 Determination of nucleotide sequence of plasmid containing
isolated human SLC-1 cDNA

After the plasmid obtained in Reference Example 1-
5 was introduced into ELECTROMAX™ DH10B™ Cells by the
electroporation method, clones with cDNA inserted
30 fragments were selected in LB agar culture medium
containing ampicillin and X-gal. Using a sterilized
toothpick, only the clones showing white color were
separated to give transformant E. coli DH10B/hSLC-1.
Individual clones were cultured overnight in LB culture
35 medium containing ampicillin, and the plasmid DNA was
refined using QIA prep8 mini prep (Qiagen). The reactions

to determine nucleotide sequence were conducted using a DyeDeoxy Terminator Cycle Sequence Kit (Parkin Elmer Co.), and the nucleotide sequence was decoded using a fluorescent light automatic sequencer.

5 As the results, obtained was the sequence shown in Sequence number: 8. The amino acid sequence (Sequence number: 9) coded by the nucleotide sequence obtained here, differs from the human SLC-1 amino acid sequence predicted as the sequence analogized from rat SLC-1 based on human
10 chromosome DNA sequence (accession number: Z86090) containing human SLC-1 sequence, in the report by Lakaye, et al. (Lakaye, B., et al. (1998) Biochim. Biophys. Acta. Vol. 1401, pp. 216-220). This shows the presence of ATG, the initiation codon, on mRNA, in the 69 and 64 amino acids
15 upstream from the estimated sequence. Escherichia coli DH10B/phSLC1L8, the transformant produced by the plasmid containing DNA coding this sequence was deposited at IFO and NIBH.

20 Reference Example 1-7

Amplification of human SLC-1cDNA by PCR method using human fetal brain originated cDNA

Amplification by the PCR method was conducted using as the template plasmid containing human SLC-1 DNA sequence
25 cloned by the gene trap method, and using synthetic DNA primers of sequence number: 10 and sequence number: 11, and synthetic DNA primers of sequence number: 12 and sequence number: 13, respectively. The former amplified DNA and the latter amplified DNA were named as "human SLC-1(S)" and
30 "human SLC-1(L)", respectively. The synthetic DNA primer was constructed so that the genes in the domain translated to the receptor protein were amplified. At that time, a recognition sequence for each restriction enzyme was added on the 5' side and 3' side, so that the nucleotide sequence
35 recognized by restriction enzyme Sal I would be added on the 5' side of the gene, and the nucleotide sequence

WO 01/21577

PCT/JP00/06375

330

recognized by restriction enzyme Spe I would be added on the 3' side. The composition of the reaction mixture for human SLC-1(S) amplification was: 5 µl of plasmid template containing human SLC-1 DNA sequence, 0.4 µM of respective synthetic DNA primers, 0.2 mM of dNTPs and 0.5 µl of Pfu DNA polymerase and buffers attached to the enzyme, with total quantity for reaction set at 50 µl. A thermal cycler (Parkin Elmer Co.) was used for the cycles for amplification. After heating at 94°C for 60 seconds, the cycle consisting of 94°C for 60 seconds, 57°C for 60 seconds, and 72°C for 150 seconds, was repeated 25 times, and finally the temperature of the reactant was maintained at 72°C for 10 minutes. The composition of the reaction mixture for human SLC-1(L) amplification was 5 µl of plasmid template containing human SLC-1 DNA sequence, 0.4 µM of respective synthetic DNA primers, 0.2 mM of dNTPs, 0.5 µl of Pfu DNA polymerase and buffers attached to the enzymes, with total quantity for reaction set at 50 µl. A thermal cycler (Parkin Elmer Co.) was used for the cycles for amplification. After heating at 94°C for 60 seconds, the cycle consisting of 94°C for 60 seconds, 60°C for 60 seconds, and 72°C for 3 minutes, was repeated 25 times, and finally the temperature of the reactant was maintained at 72°C for 10 minutes. After 0.8% agarose gel electrophoresis, confirmation of amplified products was conducted by ethidium bromide dying.

Reference Example 1-8

Subcloning of PCR product into plasmid vector and confirmation of amplified cDNA sequence by decoding of nucleotide sequence of inserted cDNA section

The reaction product after PCR in Reference Example 1-7 was separated using 0.8% low-melting point agarose gel, and the band section was cut out using a razor. After that, fragmentation, phenol extraction, phenol-chloroform extraction, and ethanol precipitation were conducted, and

the DNA was recovered. The recovered DNA was subcloned into pCR-Script Amp SK(+) plasmid vector, as prescribed by the PCR-Script™ Amp SK(+) cloning kit (Stratagene Co.). After this was introduced into Escherichia coli DH5a competent
5 cells (TOYOBO) and transformed, the clones with cDNA inserted fragments were selected in LB agar culture medium containing ampicillin and X-gal. Using a sterilized toothpick, only clones showing white color were separated to give E. coli DH5α/hSLC-1(S), which is a transformant of
10 human SLC-1 (S), and E. coli DH5α/hSLC-1(L), which is a transformant of human SLC-1 (L). Each clone was cultured overnight in LB culture medium containing ampicillin, and plasmid DNA was prepared using QIA prep8 mini prep (Qiagen).

Some of the prepared DNA was digested with Sal I and Spe
15 I restriction enzymes, and the size of the receptor cDNA fragments inserted was confirmed. The reactions to determine nucleotide sequence were conducted using a DyeDeoxy Terminator Cycle Sequence Kit (Parkin Elmer Co.) and the nucleotide sequence was decoded using a fluorescent
20 light automatic sequencer. The sequence of the obtained clones respectively matched the DNA sequence (sequence number:14) which should be amplified by synthetic DNA primers of sequence number: 10 and sequence number: 11 using human SLC-1 gene as a template, and the DNA sequence
25 (sequence number: 15) which should be amplified by synthetic DNA primers of sequence number: 12 and sequence number: 13 using human SLC-1 gene as a template.

Reference Example 1-9

30 Preparation of CHO cells for expression of human SLC-1(S), and CHO cells for expression of human SLC-1(L)

Plasmid was prepared from the E. coli clones transformed by the plasmid wherein inserted were human SLC-1(S) and human SLC-1(L) whose sequences were confirmed
35 in Reference Example 1-8, using a Plasmid Midi Kit (Qiagen), and the insert section was cut out using Sal I and Spe I

restriction enzymes. After electrophoresis was conducted, the insert DNA was cut out from agarose gel using a razor. Next, fragmentation, phenol extraction, phenol-chloroform extraction, and ethanol precipitation were conducted, and the insert DNA was recovered.

This insert DNA was added to pAKKO-111H vector plasmid for animal cell expression, digested with Sal I and Spe I (the same vector plasmid as the pAKKO1.11H described in Hinuma, S., et al., Biochim. Biophys. Acta, Vol. 1219, pp. 251-259 (1994)), and ligation was conducted by adding T4 ligase (TaKaRa Shuzo), to construct pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) plasmids for protein expression.

After E. coli DH5 α (TOYOBO) transformed by pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) was cultured, pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) plasmid DNAs were prepared using a Plasmid Midi Kit (Qiagen). These were introduced into CHO dhfr⁻ cells in accordance with the attached protocol, using a CellPfect Transfection Kit (Amersham Pharmacia Biotech Co.). A coprecipitative suspension of 10 μ g of DNA with calcium phosphate was made, which was added to 10 cm Petri dishes seeded 24 hours in advance with 5×10^5 or 1×10^6 CHO dhfr⁻ cells. After the above was cultured for 1 day in MEM α culture medium containing 10% fetal bovine serum, subculture was conducted, and then cultivation was conducted in MEM α culture medium containing no nucleic acid but containing 10% dialyzed fetal bovine serum, which is a selective culture medium. 56 clones of colonies of transformed cells which are human SLC-1(S) gene introduced CHO cells, and 61 clones of colonies of transformed cells which are human SLC-1(L) gene introduced CHO cells, both of which proliferated in the selective culture medium, were selected.

Reference Example 1-10

Selection of cell colonies into which genes with large quantities of human SLC-1(S) and human SLC-1 (L) mRNA

expression have been introduced

The quantities of expressed mRNA of 56 clones of CHO/hSLC-1(S) colonies and 61 clones of CHO/hSLC-1(L) colonies, both of which were established in Reference Example 1-9, were measured in accordance with the attached protocol using a Cytostar T Plate (Amersham Pharmacia Biotech Co.) as shown below.

After each well of the Cytostar T Plate was seeded with each clone of CHO/hSLC-1(S) colonies and CHO/hSLC-1(L) colonies by 2.5×10^4 , and cultured for 24 hours, the cells were fixed using 10% formalin.

After 0.25% Triton X-100 was added to each well to increase cell permeability, ^{35}S -labeled riboprobe of sequence number: 16 was added and hybridization was conducted.

20 mg/ml of RNaseA was added to each well to digest free riboprobe. After the plate was washed well, the radioactivity of the hybridized riboprobe was determined. Colonies showing high radioactivity expressed large quantities of mRNA. Of the 7 clones which expressed large quantities of mRNA, mainly used was Clone number 57.

Experimental Example 1

Determination of antagonist activity using GTPγS binding assay of test compound

Membrane fraction was prepared by the following method, using the human SLC-1 expressing CHO cell clone 57 obtained in Reference Example 1-10, and the rat SLC-1 expressing CHO cell clone 44 obtained in Reference Example 1-4.

The human and rat SLC-1 expressing CHO cells (1×10^8) were scraped in buffer saline phosphate (pH 7.4) to which 5 mM EDTA (ethylenediaminetetraacetic acid) had been added, and centrifuged. 10 ml of homogenized buffer (10 mM NaHCO_3 , 5 mM EDTA, pH 7.5) was added to the cell pellets, and they were homogenized using a Polytron homogenizer. The

supernatant obtained by centrifugation at $400 \times g$ for 15 minutes was further centrifuged at $100,000 \times g$ for 1 hour, to obtain the membrane fraction precipitate. This precipitate was suspended in 2 ml of assay buffer [50 mM Tris-HCl(pH 7.5), 1 mM EDTA, 0.1% BSA (bovine serum albumin), 10 mM $MgCl_2$, 100 mM NaCl, 1 μ M GDP (guanosine 5'-diphosphate), 0.25 mM PMSF (phenylmethylsulfonyl fluoride), 1 mg/ml pepstatin, 20 mg/ml leupeptin, 10 mg/ml phosphoramidon], which was centrifuged at $100,000 \times g$ for 1 hour. The membrane fraction recovered as precipitate was suspended again in 2 ml of assay buffer, and after the suspension was divided, individual portions were preserved at $-80^\circ C$ and thawed before every use.

Determination of antagonist activity of the test compound was conducted as shown below. After 171 μ l of SLC-1 expressing CHO cell membrane fractions diluted with assay buffer was poured into each well of a 96-well polypropylene plate, 2 μ l of 3×10^{-10} M MCH diluted with DMSO solution, 2 μ l of test compound solution diluted to various concentrations, and 25 μ l of [^{35}S]-Guanosine 5'-(γ -thio) triphosphate (produced by Daiichi Kagaku Yakuhin) were added respectively. (Final concentration of cell membrane: 20 μ g/ml, final concentration of [^{35}S]-Guanosine 5'-(γ -thio) triphosphate: 0.33 nM).

After this reaction mixture was allowed to react for 1 hour under stirring, it was filtered under vacuum using a glass filter (GF-C), then the filter was washed 3 times with 300 μ l of washing solution (50 mM Tris-HCl buffer solution pH 7.5). 50 ml of liquid scintillator was added to the glass filter, and residual radioactivity was determined using a liquid scintillation counter.

The IC_{50} value of the compound was calculated from the binding inhibition rate (%), based on the definition that the binding inhibition rate (%) = (radioactivity when compound and MCH were added - radioactivity when DMSO solution was added)/(radioactivity when MCH was added -

WO 01/21577

PCT/JP00/06375

335

radioactivity when DMSO solution was added) $\times 100$.

The results were shown below.

Compound Number	Inhibition Activity (IC ₅₀ value: nM)
Reference Example 25	90
Example 1	40

5

Industrial Applicability

Compounds (I), (I') and salts thereof possess excellent MCH receptor antagonistic activities, and are useful as an agent for preventing or treating obesity, etc.

10


WO 01/21577

PCT/JP00/06375

336

CLAIMS

1. A melanin-concentrating hormone antagonist which comprises a compound of the formula :



5 wherein Ar¹ is a cyclic group which may have substituents;
X is a spacer having a main chain of 1 to 6 atoms;
Y is a bond or a spacer having a main chain of 1 to 6 atoms;
Ar is a monocyclic aromatic ring which may be condensed with
10 a 4 to 8 membered non-aromatic ring, and may have further substituents;

R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing
15 hetero ring which may have substituents; R² may form a spiro ring together with Ar; or R², together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents; or a salt thereof.

20 2. An antagonist according to claim 1, wherein Y is a spacer having a main chain of 1 to 6 atoms; R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring
25 which may have substituents; or R² may form a spiro ring together with Ar.

3. An antagonist according to claim 2, wherein Ar¹ is an aromatic group which may have substituents; and "a hydrocarbon group which may have substituents" for R¹ and
30 R² is "C₁₋₆ alkyl which may have substituents".

4. An antagonist according to claim 1, wherein the cyclic

group for Ar¹ is C₆₋₁₄ monocyclic or condensed polycyclic aromatic hydrocarbon group.

- 5 5. An antagonist according to claim 1, wherein the cyclic group for Ar¹ is a group formed by removing an optional one hydrogen atom from an aromatic ring assemble in which 2 or 3 C₆₋₁₄ monocyclic or condensed polycyclic aromatic hydrocarbon groups are directly bonded by single bonds.
- 10 6. An antagonist according to claim 1, wherein the cyclic group for Ar¹ is a group formed by removing an optional one hydrogen atom from an aromatic ring assemble in which C₆₋₁₄ monocyclic or condensed polycyclic aromatic hydrocarbon and 5 to 10 membered aromatic hetero ring are directly
15 bonded by a single bond.

7. An antagonist according to claim 1, wherein Ar¹ is phenyl, biphenyl, phenyl-pyridyl, phenyl-furyl, phenyl-isoxazolyl, diphenyl-oxazolyl, pyridyl-phenyl,
20 phenyl-pyrimidinyl, benzofuranyl-phenyl, furyl-phenyl, terphenyl, thienyl-phenyl, indolyl, naphthyl-oxadiazolyl, benzofuranyl-oxadiazolyl, benzothienyl, benzofuranyl, fluorenyl, pyridyl-pyrrolyl or thioxanthanyl;
- 25 each of which may have 1 to 3 substituents selected from the group consisting of halogen atom; nitro; C₁₋₃ alkylenedioxy; optionally halogenated C₁₋₆ alkyl; hydroxy-C₁₋₆ alkyl; optionally halogenated C₃₋₆ cycloalkyl; optionally halogenated C₁₋₆ alkoxy; optionally halogenated
30 C₁₋₆ alkythio; hydroxy; C₇₋₁₉ aralkyloxy which may have substituents; C₆₋₁₄ aryloxy which may have substituents; amino; mono-C₁₋₆ alkylamino; di-C₁₋₆ alkylamino; 5 to 7 membered saturated cyclic amino which may have substituents and may be condensed with a benzene ring; 5 to 7 membered
35 non-aromatic heterocyclic groups which may have substituents; formyl; carboxy; C₆₋₁₄ aryl-carbonyl which may

have substituents; C₆₋₁₄ aryl-carbamoyl which may have substituents; aromatic hetero ring-carbamoyl which may have substituents; C₁₋₆ alkoxy-carbonyl; optionally halogenated C₁₋₆ alkyl-carboxamide; C₆₋₁₄ aryl-carboxamide
5 which may have substituents; C₇₋₁₉ aralkyl-carboxamide which may have substituents; aromatic hetero ring-carboxamide which may have substituents; N-(C₆₋₁₄ aryl-carbonyl which may have substituents)-N-C₁₋₆ alkylamino; C₆₋₁₄ arylamino-carbonylamino which may have substituents; C₆₋₁₄
10 arylsulfonylamino which may have substituents; C₆₋₁₄ aryl-carbonyloxy which may have substituents; oxo; carboxy-C₁₋₆ alkyl; C₁₋₆ alkoxy-carbonyl-C₁₋₆ alkyl; C₇₋₁₉ aralkyl which may have substituents; aromatic hetero ring-C₁₋₆ alkoxy; and cyano.

15
8. An antagonist according to claim 1, wherein Ar¹ is piperidinyl, piperazinyl, pyrrolidinyl, dihydropyridyl or tetrahydropyridyl; each of which may have 1 or 2 substituents selected from the group consisting of oxo, C₆₋₁₄
20 aryl which may have substituents, hydroxy, C₇₋₁₉ aralkyloxy-carbonyl, and C₇₋₁₉ aralkyl.


9. An antagonist according to claim 1, wherein the "spacer having a main chain of 1 to 6 atoms" for X and Y
25 is a bivalent group consisting of 1 to 3 species selected from -O-, -S-, -CO-, -SO-, -SO₂-, -NR⁸- (R⁸ is hydrogen atom, optionally halogenated C₁₋₆ alkyl, optionally halogenated C₁₋₆ alkyl-carbonyl, optionally halogenated C₁₋₆ alkylsulfonyl), and a bivalent C₁₋₆ non-cyclic hydrocarbon
30 group which may have substituents.

10. An antagonist according to claim 1, wherein X is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}-
wherein R^{8c} is hydrogen atom or C₁₋₆ alkyl.

35
11. An antagonist according to claim 1, wherein Y is an

optionally halogenated bivalent C_{1-6} non-cyclic hydrocarbon group.

12. An antagonist according to claim 1, wherein Ar is a ring of the formula :




wherein ----- is a single bond or double bond, n is an integer of 1 to 4.

13. An antagonist according to claim 1, wherein R^1 and R^2 are hydrogen atom or C_{1-6} alkyl which may have substituents; or R^1 and R^2 , together with the adjacent nitrogen atom, form a 3 to 8 membered nitrogen-containing hetero ring.
14. An antagonist according to claim 1, which is an agent for preventing or treating diseases caused by a melanin-concentrating hormone.
15. An antagonist according to claim 1, which is an agent for preventing or treating obesity.
16. An antagonist according to claim 1, which is an anorectic agent.
17. A pharmaceutical, which comprises a melanin-concentrating hormone antagonist in combination with at least one species selected from the group consisting of an agent for treating diabetes, an agent for treating hypertension and an agent for treating arteriosclerosis.
18. A compound of the formula :


WO 01/21577

PCT/JP00/06375

340




wherein Ar^1 is a cyclic group which may have substituents;
 Ar' is a ring of the formula :



- 5 wherein ----- is a single bond or double bond, n is an integer of 1 to 4, and each ring may have substituents;
 X' is $-\text{CONR}^{\text{Bc}}-$, $-\text{NR}^{\text{Bc}}\text{CO}-$, $-\text{CH}=\text{CH}-\text{CONR}^{\text{Bc}}-$ or $-\text{SO}_2\text{NR}^{\text{Bc}}-$ where R^{Bc} is hydrogen atom or C_{1-6} alkyl;
 Y is a spacer having a main chain of 1 to 6 atoms;
10 R^1 and R^2 are independently hydrogen atom or a hydrocarbon group which may have substituents; R^1 and R^2 , together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R^2 , together with the adjacent nitrogen atom and Y , may form a
15 nitrogen-containing hetero ring which may have substituents;

provided that Ar' is a ring of the formula :




- wherein symbols have the same meanings as defined above,
20 and each ring may have substituents, when X' is $-\text{SO}_2\text{NH}-$;
and provided that Ar^1 is not biphenyl which may be substituted, when X' is $-\text{CONH}-$ and Ar' is any one of benzopyran, dihydrobenzopyran, dihydrobenzoxazine, dihydrobenzoxazole or tetrahydrobenzoxazepine;
25 (excluding $\text{N}-[2-(\text{N},\text{N}-\text{dimethylamino})\text{methyl}-6\text{-tetralinyl}]-4\text{-biphenylcarboxamide}$); or a salt thereof.

19. A compound of the formula :

WO 01/21577

PCT/JP00/06375

341



wherein Ar¹ is a cyclic group which may have substituents;

----- is a single bond or double bond;

n is an integer of 1 to 4;


5 X' is -CONR^{8c}-, -NR^{8c}CO- or -CH=CH-CONR^{8c}- where R^{8c} is hydrogen atom or C₁₋₆ alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;

R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with

10 the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R², together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;


15 a ring of the formula :



wherein symbols have the same meanings as defined above, may have further substituents;

20 provided that N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]-4-biphenylcarboxamide is excluded; or a salt thereof.

20. A compound according to claim 19, which is of the formula :



25

wherein R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R²,

WO 01/21577

PCT/JP00/06375

342


together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; the other symbols have the same meanings as defined in claim 19.

5

21. A compound according to claim 20, wherein Ar¹ is an aromatic group which may have substituents; and "a hydrocarbon group which may have substituents" for R¹ and R² is "C₁₋₆ alkyl which may have substituents".

10

22. A compound of the formula :




wherein Ar¹ is a cyclic group which may have substituents; n is an integer of 1 to 4;

15 X' is -CONR^{8c}-, -NR^{8c}CO- or -CH=CH-CONR^{8c}- where R^{8c} is hydrogen atom or C₁₋₆ alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;

R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R², together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;

25 a ring of the formula :



wherein n has the same meaning as defined above, may have further substituents;

provided that N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]-4-biphenylcarboxamide is excluded; or a salt

30


WO 01/21577

PCT/JP00/06375

343

thereof.

23. A compound according to claim 22, which is of the formula :




wherein R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; the other symbols have the same meanings as defined in claim 22.

10

24. A compound according to claim 23, wherein Ar¹ is an aromatic group which may have substituents; and "a hydrocarbon group which may have substituents" for R¹ and R² is "C₁-₆ alkyl which may have substituents".

15

25. A compound of the formula :



wherein Ar¹ is a cyclic group which may have substituents; X' is -CONR^{8c}-, -NR^{8c}CO- or -CH=CH-CONR^{8c}- where R^{8c} is hydrogen atom or C₁-₆ alkyl; Y is a spacer having a main chain of 1 to 6 atoms; R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; or R², together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;

20

25


30

WO 01/21577

PCT/JP00/06375


344

a ring of the formula :



may have further substituents; or a salt thereof.


- 5 26. A compound according to claim 25, which is of the formula :



- 10 wherein R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; the other symbols have the same meanings as defined in claim 25.

- 15 27. A compound according to claim 26, wherein Ar¹ is an aromatic group which may have substituents; and "a hydrocarbon group which may have substituents" for R¹ and R² is "C₁-₆ alkyl which may have substituents".

- 20 28. A compound of the formula :



- 25 wherein Ar¹ is a cyclic group which may have substituents; X' is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}- where R^{8c} is hydrogen atom or C₁-₆ alkyl; Y is a spacer having a main chain of 1 to 6 atoms; R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with the adjacent nitrogen atom, may form a nitrogen-containing


WO 01/21577

PCT/JP00/06375

345

hetero ring which may have substituents; or R^2 , together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;


5 a ring of the formula :



may have further substituents;
provided that Ar^1 is not biphenyl which may be substituted, when X' is $-CONH-$; or a salt thereof.

10


29. A compound of the formula :



wherein Ar^1 is a cyclic group which may have substituents;
 X' is $-CONR^{8c}-$, $-NR^{8c}CO-$, $-CH=CH-CONR^{8c}-$ or $-SO_2NR^{8c}-$ where
15 R^{8c} is hydrogen atom or C_{1-6} alkyl;

Y is a spacer having a main chain of 1 to 6 atoms;
 R^1 and R^2 are independently hydrogen atom or a hydrocarbon group which may have substituents; R^1 and R^2 , together with the adjacent nitrogen atom, may form a nitrogen-containing
20 hetero ring which may have substituents; or R^2 , together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents;

a ring of the formula :



25


may have further substituents; or a salt thereof.

30. A compound of the formula :

WO 01/21577

PCT/JP00/06375


346



wherein Ar¹ is a cyclic group which may have substituents;
X' is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}- where
R^{8c} is hydrogen atom or C₁₋₆ alkyl;


- 5 Y is a spacer having a main chain of 1 to 6 atoms;
R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; or R², together
10 with the adjacent nitrogen atom and Y, may form a
nitrogen-containing hetero ring which may have
substituents;

a ring of the formula :



- 15 may have further substituents; or a salt thereof.

31. A compound of the formula :



- wherein Ar¹ is a cyclic group which may have substituents;
20 X' is -CONR^{8c}-, -NR^{8c}CO-, -CH=CH-CONR^{8c}- or -SO₂NR^{8c}- where
R^{8c} is hydrogen atom or C₁₋₆ alkyl;
Y is a spacer having a main chain of 1 to 6 atoms;
R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
25 the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; or R², together
with the adjacent nitrogen atom and Y, may form a
nitrogen-containing hetero ring which may have


WO 01/21577

PCT/JP00/06375

347

substituents;

a ring of the formula :



may have further substituents;

5 provided that Ar¹ is not biphenyl which may be substituted, when X' is -CONH-; or a salt thereof.

32. A pharmaceutical composition which comprises a compound as defined in any one of claims 18, 19, 22, 25,
10 26, 28, 29, 30 and 31.


33. A prodrug of a compound as defined in any one of claims 18, 19, 22, 25, 26, 28, 29, 30 and 31.

15 34. A compound according to claim 18, which is
N-[2-(N,N-dimethylamino)methyl-6-tetralinyl]-(4'-methoxybiphenyl-4-yl)carboxamide;
4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
20 4'-fluoro-N-[6-(1-piperidinylmethyl)-7,8-dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
25 (+)-4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
(-)-4'-fluoro-N-[6-[(N,N-dimethylamino)methyl]-5,6,7,8-tetrahydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
30 4'-chloro-N-[3-[(N,N-dimethylamino)methyl]-2H-chromen-7-yl][1,1'-biphenyl]-4-carboxamide;
4'-fluoro-N-[6-(1-pyrrolidinylmethyl)-7,8-dihydro-2-

- naphthalenyl][1,1'-biphenyl]-4-carboxamide;
N-[3-[(dimethylamino)methyl]-2H-chromen-7-yl]-4'-
fluoro[1,1'-biphenyl]-4-carboxamide;
4'-chloro-N-[6-[(dimethylamino)methyl]-5-methyl-7,8-
5 dihydro-2-naphthalenyl][1,1'-biphenyl]-4-carboxamide;
6-(4-methoxyphenyl)-N-[5-methyl-6-(1-
pyrrolidinylmethyl)-7,8-dihydro-2-
naphthalenyl]nicotinamide;
4'-chloro-N-[7-[(dimethylamino)methyl]-5,6-dihydro-3-
10 quinoliny][1,1'-biphenyl]-4-carboxamide;
4-(4-chlorophenyl)-N-[6-(1-pyrrolidinylmethyl)-7,8-
dihydro-2-naphthalenyl]-3,6-dihydro-1(2H)-
pyridinecarboxamide;
N-[6-[(dimethylamino)methyl]-7,8-dihydro-2-
15 naphthalenyl]-4-(4-fluorophenyl)-1-
piperidinecarboxamide;
4-(4-methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-5-
methyl-7,8-dihydro-2-naphthalenyl]-1-
piperidinecarboxamide;
20 4'-fluoro-N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-
naphthalenyl][1,1'-biphenyl]-4-carboxamide;
4'-chloro-N-[6-[2-(1-pyrrolidinyl)ethyl]-7,8-dihydro-2-
naphthalenyl][1,1'-biphenyl]-4-carboxamide;
4'-chloro-N-[2-[(dimethylamino)methyl]-3,4-dihydro-2H-
25 1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide;
4-(4-methoxyphenyl)-N-[5-methyl-6-(1-
pyrrolidinylmethyl)-7,8-dihydro-2-naphthalenyl]-1-
piperidinecarboxamide;
4-(4-chlorophenyl)-N-[6-[(4-methyl-1-
30 piperazinyl)methyl]-7,8-dihydro-2-naphthalenyl]-1-
piperidinecarboxamide;
4'-chloro-N-[2-[(dimethylamino)methyl]-1H-inden-6-
yl][1,1'-biphenyl]-4-carboxamide;
4'-fluoro-N-[2-(1-pyrrolidinylmethyl)-3,4-dihydro-2H-
35 1,4-benzoxazin-6-yl][1,1'-biphenyl]-4-carboxamide;
4'-fluoro-N-[5-methyl-6-[(4-methyl-1-

piperaziny]methyl]-7,8-dihydro-2-naphthalenyl][1,1'-
biphenyl]-4-carboxamide;
4'-chloro-N-[5-methyl-6-[(4-methyl-1-
piperaziny]methyl]-7,8-dihydro-2-naphthalenyl][1,1'-
5 biphenyl]-4-carboxamide; or
4-(4-chlorophenyl)-N-[5-methyl-6-[(4-methyl-1-
piperaziny]methyl]-7,8-dihydro-2-naphthalenyl]-1-
piperidinecarboxamide.

- 10 35. A method for preventing or treating diseases caused
by a melanin-concentrating hormone in a mammal in need
thereof, which comprises administering to said mammal an
effective amount of a compound of the formula :




- 15 wherein Ar¹ is a cyclic group which may have substituents;
X is a spacer having a main chain of 1 to 6 atoms;
Y is a bond or a spacer having a main chain of 1 to 6 atoms;
Ar is a monocyclic aromatic ring which may be condensed with
a 4 to 8 membered non-aromatic ring, and may have further
20 substituents;
R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; R² may form a spiro
25 ring together with Ar; or R², together with the adjacent
nitrogen atom and Y, may form a nitrogen-containing hetero
ring which may have substituents; or a salt thereof.

36. A method for preventing or treating obesity in a mammal
30 in need thereof, which comprises administering to said
mammal an effective amount of a compound of the formula :

WO 01/21577

PCT/JP00/06375

350




wherein Ar¹ is a cyclic group which may have substituents;
X is a spacer having a main chain of 1 to 6 atoms;

Y is a bond or a spacer having a main chain of 1 to 6 atoms;
Ar is a monocyclic aromatic ring which may be condensed with
a 4 to 8 membered non-aromatic ring, and may have further
substituents;

R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; R² may form a spiro
ring together with Ar; or R², together with the adjacent
nitrogen atom and Y, may form a nitrogen-containing hetero
ring which may have substituents; or a salt thereof.

37. Use of a compound of the formula :




wherein Ar¹ is a cyclic group which may have substituents;
X is a spacer having a main chain of 1 to 6 atoms;

Y is a bond or a spacer having a main chain of 1 to 6 atoms;
Ar is a monocyclic aromatic ring which may be condensed with
a 4 to 8 membered non-aromatic ring, and may have further
substituents;

R¹ and R² are independently hydrogen atom or a hydrocarbon
group which may have substituents; R¹ and R², together with
the adjacent nitrogen atom, may form a nitrogen-containing
hetero ring which may have substituents; R² may form a spiro
ring together with Ar; or R², together with the adjacent
nitrogen atom and Y, may form a nitrogen-containing hetero
ring which may have substituents; or a salt thereof;

for the manufacture of a pharmaceutical preparation for preventing or treating diseases caused by a melanin-concentrating hormone.

5 38. Use of a compound of the formula :



wherein Ar¹ is a cyclic group which may have substituents;
X is a spacer having a main chain of 1 to 6 atoms;
Y is a bond or a spacer having a main chain of 1 to 6 atoms;
10 Ar is a monocyclic aromatic ring which may be condensed with a 4 to 8 membered non-aromatic ring, and may have further substituents;
R¹ and R² are independently hydrogen atom or a hydrocarbon group which may have substituents; R¹ and R², together with
15 the adjacent nitrogen atom, may form a nitrogen-containing hetero ring which may have substituents; R² may form a spiro ring together with Ar; or R², together with the adjacent nitrogen atom and Y, may form a nitrogen-containing hetero ring which may have substituents; or a salt thereof;
20 for the manufacture of a pharmaceutical preparation for preventing or treating obesity.

WO 01/21577

PCT/JP00/06375

1/11

SEQUENCE LISTING

<110> Takeda Chemical Industries, Ltd.
 <120> Melanin Concentrating Hormone Antagonist
 <130> 2648W00P
 <150> JP 11-266298
 <151> 1999-09-20
 <150> JP 11-357889
 <151> 1999-12-16
 <150> JP 2000-126272
 <151> 2000-04-20
 <160> 16
 <210> 1
 <211> 32
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223>
 <400> 1
 GTCGACATGG ATCTGCAAAC CTCGTTGCTG TG 32
 <210> 2
 <211> 32
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223>
 <400> 2
 ACTAGTTCAG GTGCCTTTGC TTTCTGTCCT CT 32
 <210> 3

WO 01/21577

PCT/JP00/06375

2/11

<211> 353

<212> PRT

<213> Rat

<400> 3

```

Met Asp Leu Gln Thr Ser Leu Leu Ser Thr Gly Pro Asn Ala Ser Asn
1           5           10           15
Ile Ser Asp Gly Gln Asp Asn Leu Thr Leu Pro Gly Ser Pro Pro Arg
           20           25           30
Thr Gly Ser Val Ser Tyr Ile Asn Ile Ile Met Pro Ser Val Phe Gly
           35           40           45
Thr Ile Cys Leu Leu Gly Ile Val Gly Asn Ser Thr Val Ile Phe Ala
           50           55           60
Val Val Lys Lys Ser Lys Leu His Trp Cys Ser Asn Val Pro Asp Ile
65           70           75           80
Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu Phe Leu Leu Gly Met
           85           90           95
Pro Phe Met Ile His Gln Leu Met Gly Asn Gly Val Trp His Phe Gly
           100          105          110
Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp Ala Asn Ser Gln Phe
           115          120          125
Thr Ser Thr Tyr Ile Leu Thr Ala Met Thr Ile Asp Arg Tyr Leu Ala
           130          135          140
Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg Lys Pro Ser Met Ala
145          150          155          160
Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr
           165          170          175
Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val
           180          185          190
Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe

```

WO 01/21577

PCT/JP00/06375

3/11

195	200	205
Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile		
210	215	220
Thr Ala Ala Tyr Val Lys Ile Leu Gln Arg Met Thr Ser Ser Val Ala		
225	230	235
Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg		
245	250	255
Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr		
260	265	270
Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr		
275	280	285
Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser		
290	295	300
Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Thr Phe Arg Lys		
305	310	315
Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln Gly Gln Leu Arg Thr		
325	330	335
Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg Thr Glu Ser Lys Gly		
340	345	350

Thr

<210> 4

<211> 1074

<212> DNA

<213> Rat

<400> 4

GTCGACATGG ATCTGCAAAC CTCGTTGCTG TCCACTGGCC CCAATGCCAG CAACATCTCC	60
GATGGCCAGG ATAATCTCAC ATTGCCGGGG TCACCTCCTC GCACAGGGAG TGTCTCCTAC	120
ATCAACATCA TTATGCCTTC CGTGTTTGGT ACCATCTGTC TCCTGGGCAT CGTGGGAAAC	180
TCCACGGTCA TCITTGCTGT GGTGAAGAAG TCCAAGCTAC ACTGGTGCAG CAACGTCCCC	240

WO 01/21577

PCT/JP00/06375

4/11

GACATCTTCA TCATCAACCT CTCTGTGGTG GATCTGCTCT TCCTGCTGGG CATGCCTTTC 300
 ATGATCCACC AGCTCATGGG GAACGGCGTC TGGCACTTTG GGGAAACCAT GTGCACCCTC 360
 ATCACAGCCA TGGACGCCAA CAGTCAGTTC ACTAGCACCT ACATCCTGAC TGCCATGACC 420
 ATTGACCGCT ACTTGGCCAC CGTCCACCCC ATCTCCTCCA CCAAGTCCG GAAGCCCTCC 480
 ATGGCCACCC TGGTGATCTG CCTCCTGTGG GCGCTCTCCT TCATCAGTAT CACCCCTGTG 540
 TGGCTCTACG CCAGGCTCAT TCCCTTCCCA GGGGGTGCTG TGGGCTGTGG CATCCGCCTG 600
 CCAAACCCGG ACACTGACCT CTACTGGTTC ACTCTGTACC AGTTTTTCCT GGCCTTTGCC 660
 CTTCCGTTTG TGGTCATTAC CGCCGCATAC GTGAAAATAC TACAGCGCAT GACGTCTTCG 720
 GTGGCCCCAG CCTCCAACG CAGCATCCGG CTTCGGACAA AGAGGGTGAC CCGCACGGCC 780
 ATTGCCATCT GTCTGGTCTT CTTTGTGTGC TGGGCACCTT ACTATGTGCT GCAGCTGACC 840
 CAGCTGTCCA TCAGCCGCCC GACCCTCACG TTTGTCTACT TGTACAACGC GGCCATCAGC 900
 TTGGGCTATG CTAACAGCTG CCTGAACCCC TTTGTGTACA TAGTGCTCTG TGAGACCTTT 960
 CGAAAACGCT TGGTGTGTC AGTGAAGCCT GCAGCCCAGG GGCAGCTCCG CACGGTCAGC 1020
 AACGCTCAGA CAGCTGATGA GGAGAGGACA GAAAGCAAAG GCACCTGAAC TAGT 1074

<210> 5

<211> 262

<212> RNA

<213> Rat

<400> 5

GCGAAUUGGG UACCGGGCCC CCCUCGAGG UCGACGGUUA CGAUAAAGCU GAUAUCGAAU 60
 UCCUGCAGCC CGGGGGAUCC GCCCACUAGU UCAGGUGCCU UUGCUUUCUG UCCUCUCCUC 120
 AUCAGCUGUC UGAGCGUUGC UGACCGUGCG GAGCUGCCCC UGGGCUGCAG GCUUCACUGA 180
 CAACACCAAG CGUUUUCGAA AGGUCUCACA GAGCACUAUG UACACAAAGG GGUUCAGGCA 240
 GCUGUUAGCA UAGCCCAAGC UG 262

<210> 6

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

WO 01/21577

PCT/JP00/06375

5/11

<223>

<400> 6

CAACAGCTGC CTCAACCC 18

<210> 7

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

<223>

<400> 7

CCTGGTGATC TGCCTCCT 18

<210> 8

<211> 1275

<212> DNA

<213> Human

<400> 8

TAGGTGATGT CAGTGGGAGC CATGAAGAAG GGAGTGGGGA GGGCAGTTGG GCTTGGAGGC 60
GGCAGCGGCT GCCAGGCTAC GGAGGAAGAC CCCCTTCCCA ACTGCGGGGC TTGCGCTCCG 120
GGACAAGGTG GCAGGCGCTG GAGGCTGCCG CAGCCTGCGT GGGTGGAGGG GAGCTCAGCT 180
CGGTTGTGGG AGCAGGCGAC CGGCACTGGC TGGATGGACC TGAAGCCTC GCTGCTGCCC 240
ACTGGTCCCA ACGCCAGCAA CACCTCTGAT GGCCCCGATA ACCTCACTTC GGCAGGATCA 300
CCTCCTCGCA CGGGGAGCAT CTCCTACATC AACATCATCA TGCCTTCGGT GTTCGGCACC 360
ATCTGCCTCC TGGGCATCAT CGGGAAGTCC ACGGTCATCT TCGCGGTCGT GAAGAAGTCC 420
AAGCTGCACT GGTGCAACAA CGTCCCCGAC ATCTTCATCA TCAACCTCTC GGTAGTAGAT 480
CTCCTCTTTC TCCTGGGCAT GCCCTTCATG ATCCACCAGC TCATGGGCAA TGGGGTGTGG 540
CACTTTGGGG AGACCATGTG CACCCTCATC ACGGCCATGG ATGCCAATAG TCAGTTCACC 600
AGCACCTACA TCCTGACCGC CATGGCCATT GACCGCTACC TGGCCACTGT CCACCCCATC 660
TCTTCCACGA AGTTCCGGAA GCCCTCTGTG GCCACCCTGG TGATCTGCCT CCTGTGGGCC 720
CTCTCCTTCA TCAGCATCAC CCCTGTGTGG CTGTATGCCA GACTCATCCC CTTCACAGGA 780

WO 01/21577

PCT/JP00/06375

6/11

```

GGTGCAGTGG GCTGCGGCAT ACGCCTGCCC AACCCAGACA CTGACCTCTA CTGGTTCACC 840
CTGTACCAGT TTTTCCTGGC CTTTGCCCTG CCTTTTGTGG TCATCACAGC CGCATACGTG 900
AGGATCCTGC AGCGCATGAC GTCCTCAGTG GCCCCCGCCT CCCAGCGCAG CATCCGGGTG 960
CGGACAAAGA GGGTGACCCG CACAGCCATC GCCATCTGTC TGGTCTTCTT TGTGTGCTGG 1020
GCACCCTACT ATGTGCTACA GCTGACCCAG TTGTCCATCA GCCGCCCCGAC CCTCACCTTT 1080
GTCTACTTAT ACAATGCGGC CATCAGCTTG GGCTATGCCA ACAGCTGCCT CAACCCCTTT 1140
GTGTACATCG TGCTCTGTGA GACGTTCCGC AAACGCTTGG TCCTGTCGGT GAAGCCTGCA 1200
GCCCAGGGGC AGCTTCGCGC TGTCAGCAAC GCTCAGACGG CTGACGAGGA GAGGACAGAA 1260
AGCAAAGGCA CCTGA 1275

```

<210> 9

<211> 422

<212> PRT

<213> Human

<400> 9

```

MeT Ser Val Gly Ala MeT Lys Lys Gly Val Gly Arg Ala Val Gly Leu
1           5           10           15
Gly Gly Gly Ser Gly Cys Gln Ala Thr Glu Glu Asp Pro Leu Pro Asn
           20           25           30
Cys Gly Ala Cys Ala Pro Gly Gln Gly Gly Arg Arg Trp Arg Leu Pro
           35           40           45
Gln Pro Ala Trp Val Glu Gly Ser Ser Ala Arg Leu Trp Glu Gln Ala
           50           55           60
Thr Gly Thr Gly Trp MeT Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly
65           70           75           80
Pro Asn Ala Ser Asn Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala
           85           90           95
Gly Ser Pro Pro Arg Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile MeT
           100          105          110
Pro Ser Val Phe Gly Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser

```

WO 01/21577

PCT/JP00/06375

7/11

115	120	125	
Thr Val Ile Phe Ala Val	Val Lys Lys Ser Lys	Leu His Trp Cys Asn	
130	135	140	
Asn Val Pro Asp Ile Phe	Ile Ile Asn Leu Ser Val	Val Asp Leu Leu	
145	150	155	160
Phe Leu Leu Gly MeT Pro	Phe MeT Ile His Gln	Leu MeT Gly Asn Gly	
165	170	175	
Val Trp His Phe Gly Glu	Thr MeT Cys Thr Leu	Ile Thr Ala MeT Asp	
180	185	190	
Ala Asn Ser Gln Phe Thr	Ser Thr Tyr Ile Leu	Thr Ala MeT Ala Ile	
195	200	205	
Asp Arg Tyr Leu Ala Thr	Val His Pro Ile Ser	Ser Thr Lys Phe Arg	
210	215	220	
Lys Pro Ser Val Ala Thr	Leu Val Ile Cys Leu	Leu Trp Ala Leu Ser	
225	230	235	240
Phe Ile Ser Ile Thr Pro	Val Trp Leu Tyr Ala	Arg Leu Ile Pro Phe	
245	250	255	
Pro Gly Gly Ala Val Gly	Cys Gly Ile Arg Leu	Pro Asn Pro Asp Thr	
260	265	270	
Asp Leu Tyr Trp Phe Thr	Leu Tyr Gln Phe Phe	Leu Ala Phe Ala Leu	
275	280	285	
Pro Phe Val Val Ile Thr	Ala Ala Tyr Val Arg	Ile Leu Gln Arg MeT	
290	295	300	
Thr Ser Ser Val Ala Pro	Ala Ser Gln Arg Ser	Ile Arg Leu Arg Thr	
305	310	315	320
Lys Arg Val Thr Arg Thr	Ala Ile Ala Ile Cys	Leu Val Phe Phe Val	
325	330	335	
Cys Trp Ala Pro Tyr Tyr	Val Leu Gln Leu Thr	Gln Leu Ser Ile Ser	
340	345	350	

WO 01/21577

PCT/JP00/06375

8/11

Arg Pro Thr Leu Thr Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu

355

360

365

Gly Tyr Ala Asn Ser Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys

370

375

380

Glu Thr Phe Arg Lys Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln

385

390

395

400

Gly Gln Leu Arg Ala Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg

405

410

415

Thr Glu Ser Lys Gly Thr

420

<210> 10

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223>

<400> 10

GTCGACATGG ACCTGGAAGC CTCGCTGCTG C 31

<210> 11

<211> 31

<212> DNA

<213> Artificial Sequence

<220>

<223>

<400> 11

ACTAGTTCAG GTGCCTTTGC TTTCTGTCCT C 31

<210> 12

<211> 33

<212> DNA

WO 01/21577

PCT/JP00/06375

9/11

<213> Artificial Sequence

<220>

<223>

<400> 12

AGTCGACATG TCAGTGGGAG CCATGAAGAA GGG 33

<210> 13

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223>

<400> 13

AACTAGTTCA GGTGCCTTTG CTTTCTGTCC TCT 33

<210> 14

<211> 1074

<212> DNA

<213> Human

<400> 14

GTCGACATGG ACCTGGAAGC CTCGCTGCTG CCCACTGGTC CCAACGCCAG CAACACCTCT 60
GATGGCCCCG ATAACCTCAC TTCGGCAGGA TCACCTCCTC GCACGGGGAG CATCTCCTAC 120
ATCAACATCA TCATGCCTTC GGTGTTCCGC ACCATCTGCC TCCTGGGCAT CATCGGGAAC 180
TCCACGGTCA TCTTCGCGGT CGTGAAGAAG TCCAAGCTGC ACTGGTGCAA CAACGTCCCC 240
GACATCTTCA TCATCAACCT CTCGGTAGTA GATCTCCTCT TTCTCCTGGG CATGCCCTTC 300
ATGATCCACC AGCTCATGGG CAATGGGGTG TGGCACTTTG GGGAGACCAT GTGCACCCTC 360
ATCACGGCCA TGGATGCCAA TAGTCAGTTC ACCAGCACCT ACATCCTGAC CGCCATGGCC 420
ATTGACCGCT ACCTGGCCAC TGTCCACCCC ATCTCTTCCA CGAAGTCCG GAAGCCCTCT 480
GTGGCCACCC TGGTGATCTG CCTCCTGTGG GCCCTCTCCT TCATCAGCAT CACCCCTGTG 540
TGGCTGTATG CCAGACTCAT CCCCTTCCCA GGAGGTGCAG TGGGCTGCGG CATACGCCTG 600
CCCAACCCAG AACTGACCT CTA CTGGTTC ACCCTGTACC AGTTTTTCCT GGCCTTTGCC 660

WO 01/21577

PCT/JP00/06375

10/11

CTGCCTTTTG TGGTCATCAC AGCCGCATAC GTGAGGATCC TGCAGCGCAT GACGTCCTCA 720
GTGGCCCCCG CCTCCCAGCG CAGCATCCGG CTGCGGACAA AGAGGGTGAC CCGCACAGCC 780
ATCGCCATCT GTCTGGTCTT CTTTGTGTGC TGGGCACCCT ACTATGTGCT ACAGCTGACC 840
CAGTTGTCCA TCAGCCGCCC GACCCTCACC TTTGTCTACT TATACAATGC GGCCATCAGC 900
TTGGGCTATG CCAACAGCTG CCTCAACCCC TTTGTGTACA TCGTGCTCTG TGAGACGTTT 960
CGCAAACGCT TGGTCCTGTC GGTGAAGCCT GCAGCCCAGG GGCAGCTTCG CGCTGTCAGC 1020
AACGCTCAGA CGGCTGACGA GGAGAGGACA GAAAGCAAAG GCACCTGAAC TAGT 1074

<210> 15

<211> 1283

<212> DNA

<213> Human

<400> 15

AGTCGACATG TCAGTGGGAG CCATGAAGAA GGGAGTGGGG AGGGCAGTTG GGCTTGGAGG 60
CGGCAGCGGC TGCCAGGCTA CGGAGGAAGA CCCCCTTCCC AACTGCGGGG CTTGCGCTCC 120
GGGACAAGGT GGCAGGCGCT GGAGGCTGCC GCAGCCTGCG TGGGTGGAGG GGAGCTCAGC 180
TCGGTTGTGG GAGCAGGCGA CCGGCACTGG CTGGATGGAC CTGGAAGCCT CGCTGCTGCC 240
CACTGGTCCC AACGCCAGCA ACACCTCTGA TGGCCCCGAT AACCTCACTT CGGCAGGATC 300
ACCTCCTCGC ACGGGGAGCA TCTCCTACAT CAACATCATC ATGCCTTCGG TGTCGGCAC 360
CATCTGCCTC CTGGGCATCA TCGGGAATC CACGGTCATC TTCGCGGTCG TGAAGAAGTC 420
CAAGCTGCAC TGGTGCAACA ACGTCCCCGA CATCTTCATC ATCAACCTCT CGGTAGTAGA 480
TCTCCTCTTT CTCCTGGGCA TGCCCTTCAT GATCCACCAG CTCATGGGCA ATGGGGTGTG 540
GCACTTTGGG GAGACCATGT GCACCCTCAT CACGGCCATG GATGCCAATA GTCAGTTCAC 600
CAGCACCTAC ATCCTGACCG CCATGGCCAT TGACCGCTAC CTGGCCACTG TCCACCCCAT 660
CTCTCCACG AAGTTCCGGA AGCCCTCTGT GGCCACCCTG GTGATCTGCC TCCTGTGGGC 720
CCTCTCCTTC ATCAGCATCA CCCCTGTGTG GCTGTATGCC AGACTCATCC CCTTCCCAGG 780
AGGTGCAGTG GGCTGCGGCA TACGCCTGCC CAACCCAGAC ACTGACCTCT ACTGGTTCAC 840
CCTGTACCAG TTTTTCCTGG CCTTTGCCCT GCCTTTTGTG GTCATCACAG CCGCATACGT 900
GAGGATCCTG CAGCGCATGA CGTCCTCAGT GGCCCCCGCC TCCCAGCGCA GCATCCGGCT 960
GCGGACAAAG AGGGTGACCC GCACAGCCAT CGCCATCTGT CTGGTCTTCT TTGTGTGCTG 1020

WO 01/21577

PCT/JP00/06375

11/11

GGCACCTAC TATGTGCTAC AGCTGACCCA GTTGTCCATC AGCCGCCCCGA CCCTCACCTT 1080
TGTCTACTTA TACAATGCGG CCATCAGCTT GGGCTATGCC AACAGCTGCC TCAACCCCTT 1140
TGTGTACATC GTGCTCTGTG AGACGTTCCG CAAACGCTTG GTCCTGTCGG TGAAGCCTGC 1200
AGCCCAGGGG CAGCTTCGGG CTGTCAGCAA CGCTCAGACG GCTGACGAGG AGAGGACAGA 1260
AAGCAAAGGC ACCTGAACTA GTT 1283

<210> 16

<211> 420

<212> RNA

<213> Human

<400> 16

CAAAAGCUGG AGCUCCACCG CGGUGGCGGC CGCUCUAGCC CACUAGUUCA GGUGCCUUUG 60
CUUUCUGUCC UCUCUCGUC AGCCGUCUGA GCGUUGCUGA CAGCGCGAAG CUGCCCCUGG 120
GCUGCAGGCU UCACCGACAG GACCAAGCGU UUGCGGAACG UCUCACAGAG CACGAUGUAC 180
ACAAAGGGGU UGAGGCAGCU GUUGGCAUAG CCAAGCUGA UGGCCGCAUU GUAUAAGUAG 240
ACAAAGGUGA GGGUCGGGCG GCUGAUGGAC AACUGGGUCA GCUGUAGCAC AUAGUAGGGU 300
GCCCAGCACA CAAAGAAGAC CAGACAGAU GCGAUGGCUG UGCGGGUCAC CCUCUUUGUC 360
CGCAGCCGGA UGCUGCGCUG GGAGGCGGGG GCCACUGAGG ACGUCAUGCG CUGCAGGAUC 420